

# BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Impact of COVID-19 pandemic on mental health among general Bangladeshi population: A cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2020-045727
Article Type:	Original research
Date Submitted by the Author:	31-Oct-2020
Complete List of Authors:	Das, Rajesh; University of Asia Pacific, Department of Pharmacy Hasan, Md. Rakib; Jahangirnagar University, Department of Pharmacy Daria, Sohel; University of Asia Pacific, Department of Pharmacy Islam, Rabiul; University of Asia Pacific,
Keywords:	COVID-19, Depression & mood disorders < PSYCHIATRY, PUBLIC HEALTH

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Title:** Impact of COVID-19 pandemic on mental health among general Bangladeshi population: A cross-sectional study

**Running Title:** COVID-19 pandemic and mental health issues

**Authors:**

Rajesh Das<sup>1</sup>, Md. Rakib Hasan<sup>2</sup>, Sohel Daria<sup>1</sup>, Md. Rabiul Islam<sup>1,3\*</sup>

**Affiliations:**

<sup>1</sup>Department of Pharmacy, University of Asia Pacific, 74/A Green Road, Farmgate, Dhaka-1215, Bangladesh

<sup>2</sup>Department of Pharmacy, Jahangirnagar University, Savar, Dhaka-1335

<sup>3</sup>Department of Clinical Pharmacy and Pharmacology, Dhaka University, Dhaka-1000

**Author's Email:**

Rajesh Das<sup>1</sup> ([rajeshdas4764@gmail.com](mailto:rajeshdas4764@gmail.com))

Md. Rakib Hasan<sup>2</sup> ([rakibju38@gmail.com](mailto:rakibju38@gmail.com))

Sohel Daria ([soheldaria.pharm@gmail.com](mailto:soheldaria.pharm@gmail.com))

Md. Rabiul Islam<sup>1\*</sup> ([robi.ayaan@gmail.com](mailto:robi.ayaan@gmail.com))

**Word Count: 5531**

*(From the introduction to conclusion excluding abstract, references, declarations, etc.)*

**\*Correspondence:**

**Md. Rabiul Islam, PhD**

Assistant Professor, Department of Pharmacy, University of Asia Pacific  
74/A Green Road, Farmgate, Dhaka-1215, Bangladesh.

Email: [robi.ayaan@gmail.com](mailto:robi.ayaan@gmail.com); Cell: +8801916031831

ORCID: <https://orcid.org/0000-0003-2820-3144>

1  
2  
3 **Title:** Impact of COVID-19 pandemic on mental health among general  
4 Bangladeshi population: A cross-sectional study  
5  
6  
7

## 8 **ABSTRACT**

9  
10 **Objectives:** Loneliness, depression, anxiety, and sleep problems are the major mental illness  
11 that significantly amplified worldwide during the coronavirus (COVID-19) pandemic. At the  
12 early stage of the outbreak, lockdown and quarantine approaches were imposed by the  
13 government of Bangladesh with potential consequences on people's daily life and their mental  
14 and physical health. Along with physical and mental health, economic status, healthcare  
15 facilities, and other lifestyle factors in Bangladesh are greatly impacted by COVID-19. Here  
16 we aimed to measure the impact of the COVID-19 pandemic on mental health among the  
17 Bangladeshi population.  
18  
19  
20  
21  
22  
23

24 **Methods:** A cross-sectional study was conducted among 672 general Bangladeshi people aged  
25 between 15 and 65 years all over the country from April 15 to May 10, 2020. After taking  
26 informed consent, we conducted an online survey measuring the socio-demographic profiles  
27 and using the UCLA loneliness scale (ULS-8), patient health questionnaire-9 (PHQ-9),  
28 generalized anxiety disorder 7-item (GAD-7) scale, Pittsburgh sleep quality index (PSQI) for  
29 assessment of loneliness, depression, anxiety, and sleep problem, respectively.  
30  
31  
32  
33  
34

35 **Results:** Estimates of loneliness, depression, anxiety, and sleep problems were 71%, 38%,  
36 64%, and 74%, respectively, at any degree of severity. Statistically significant factors  
37 associated with depression were being females, low economic status, students or unemployed,  
38 and living without a family. Factors statistically foreseeing anxiety were female gender,  
39 obesity, students, or unemployed. The loneliness was observed among people who are living  
40 without their family members. Moreover, statistically significant inter-relationships among all  
41 four mental health disorders were observed.  
42  
43  
44  
45  
46  
47

48 **Conclusions:** A large portion of respondents reported mental health problems during the  
49 COVID-19 pandemic in Bangladesh. Based on the present findings, the longitudinal  
50 evaluations are recommended to measure the actual gravity of these issues during and after the  
51 pandemic. Appropriate supportive programs and interventional approaches would help to  
52 address mental health problems in Bangladesh during this COVID-19 pandemic.  
53  
54  
55  
56

57 **Keywords:** Coronavirus disease 2019; COVID-19; Mental health; Outburst; Bangladesh  
58  
59  
60

## Article Summary

### Strengths and limitations of this study

- To the best of our knowledge, this is the first-ever study carried out in Bangladesh to evaluate the impact of COVID-19 on mental health among the general Bangladeshi population with such a large and homogeneous sample.
- The study provides valuable information about loneliness, depression, anxiety, and sleep problems of the Bangladeshi population during the COVID-19 pandemic.
- During the COVID-19 pandemic, the overall mental health issues of the Bangladeshi population might be recapitulated through the findings of the present study.
- All data were obtained from online self-reports assessments; therefore, recall bias was unavoidable and not representative for those who have no internet facilities.
- The cross-sectional nature of the study is not able to measure the impact of these mental illnesses over time and the present study didn't consider the altered lifestyle of the respondents due to the COVID-19 pandemic.

## INTRODUCTION

The coronavirus 2019 (COVID-19) disease caused by the novel coronavirus strain SARS-CoV-2 was originated from Wuhan, China, at the end of 2019 and declared as pandemic on March 11, 2020, due to its uncontrolled spreading worldwide.<sup>1</sup> After spreading, till October 9, 2020, 36,804,776 COVID-19 affected cases were found where 1,067,560 died with 4% death vs. recovery. Currently, the spreading rate is highest from any previous time as daily new cases were found approximately 3.43 lacs on October 8, 2020.<sup>2</sup> Currently, more than a hundred vaccine candidates of COVID-19 are in the different developing stage but catastrophic conditions are not improved yet due to lack of proper treatment and medications.<sup>3</sup> Though some countries were succeeded to control the outbreak, most countries-imposed lockdown to limit widespread which eventually affects not only socioeconomic conditions but also the mental health of the population regardless of their age, gender, profession, etc.<sup>4</sup> COVID-19 is usually transmitted by the breathing of infected droplets or contact with infected droplets, then affects people psychologically, mentally, economically as well as collapses the health care system in nearly 200 infected countries.<sup>5</sup> About 52.1% of people felt horrified and apprehensive during the COVID-19 breakdown where 57.8-77.9% needed extra support from family and friends.<sup>6</sup>

Major epidemic and pandemic outbursts have several negative impacts on individual and collective mental health in society.<sup>4</sup> For example, in Taiwan, the severe acute respiratory syndrome (SARS) outbreak in 2003 severely impacted the mental health of healthcare professionals working in tertiary care hospitals (Chong et al., 2004). Another study in Canada reported that healthcare providers working at hospitals during the SARS outbreak experienced long-term occupational and psychological effects.<sup>7</sup> In the USA, a study was conducted among Capitol Hill staff workers to respond successfully to disaster-related mental health after the anthrax attacks in 2001. This study reported that 55% and 27% of the respondents were suffering from a post-traumatic stress disorder and post-anthrax psychiatric disorder.<sup>8</sup> The mental health of many individuals is potentially affected by COVID-19 in many ways. In addition to COVID-19 patients, close contacts, isolated suspected cases, families and friends of patients, and healthcare providers, and general people may also experience extra mental health burdens.<sup>9</sup> Therefore, understanding the impact of the COVID-19 pandemic on an individual's mental health may help to reduce many current and future mental health issues.

Bangladesh is a densely populated county with about 164 million population which is five times denser than any other mega country.<sup>10</sup> Many factors such as population density,

1  
2  
3 cultivation land saturation, poor economic background, climate change, disease burden affect  
4 mental health of general people.<sup>11</sup> The economic backbone of the general people of Bangladesh  
5 further affected after imposing the lockdown from 26<sup>th</sup> March 2020 due to the COVID-19  
6 pandemic. In Bangladesh, the country's Institute of Epidemiology, Disease Control and  
7 Research (IEDCR) reported the first COVID-19 cases on March 8, 2020.<sup>12</sup> On October 8, 2020,  
8 total cases numbered 374,592, with 5,460 fatalities and 288,316 recoveries.<sup>13</sup> As the infection  
9 spread every corner of the country and the numbers of confirmed cases significantly increased  
10 over the last six months, mental health-related disorders may increase, particularly among  
11 susceptible people. Many hospitals weren't prepared well to treat COVID-19 patients and the  
12 testing facilities were limited at the initial time that greatly impacted the mental health of  
13 common people of Bangladesh, even the situation influenced some people's suicidal ideation  
14 who failed to cope with these extra mental burdens.<sup>14-16</sup> To date, not enough attention has been  
15 paid to reduce mental health problems due to COVID-19 rather identifying and treating new  
16 cases. This data would help to address the psychological issues that people are experiencing in  
17 Bangladesh during this COVID-19 pandemic. However, few studies have considered the  
18 mental health issues among the population of Bangladesh but extensive research on different  
19 mental parameters simultaneously was not evaluated yet.<sup>17-20</sup>

20  
21  
22 In this study, we aimed to estimate the extent to which general people of Bangladesh were  
23 prone to loneliness, depression, anxiety, and sleep problems due to the COVID-19 pandemic  
24 and to discover the relationship between the mental health problems and sociodemographic  
25 factors. Also, we explored the specific factors affecting the mental health of people and  
26 suggested the possible approaches to manage the situation during and after the pandemic.

## 27 28 29 **METHODS**

### 30 31 32 *Participants and procedure*

33  
34  
35 This cross-sectional and anonymous online survey enrolled 672 respondents (381 males, 291  
36 females) aged between 15 to 65 years from all over Bangladesh. The questionnaire was  
37 circulated in both English and Bengali versions for the proper understanding of questions.  
38 Initially, 736 respondents were given feedback about the questions but the responses from 64  
39 were excluded due to the partial information. All the respondents of this survey were of  
40 Bangladeshi ethnicity and living in Bangladesh at that time. Any Bangladeshi people aged  
41 between 15 to 65 years who understood the questions properly were included in this survey.  
42 Exclusion criteria included the history of other psychiatric disorders like delusions, mental  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 retardation, bipolar disorder, schizophrenia, personality disorder, mood-congruent or  
4 incongruent psychotic features, comorbid psychiatric illness, neurological disease, or patients  
5 with clinical evidence of dementia. Respondents having severe or acute medical illnesses, the  
6 presence of other acute or chronic diseases, history of addiction, or substance abuse were also  
7 excluded from this survey. We did not pay for participation.  
8  
9

### 10 11 12 *Estimations*

13  
14 A self-reported structured questionnaire covering the informed consent, information about  
15 socio-demographics and biophysics, and psychometric assessment scales. The questionnaire  
16 was reached to participants via mail, hand to hand, or via transports that carry emergency  
17 things. Any doubt regarding the questionnaire was solved over the phone calls or video  
18 conferences. Illiterate or older respondents were helped by their educated family members. The  
19 data were collected from 15<sup>th</sup> April 2020 to 10<sup>th</sup> May 2020 using the google survey tool (Google  
20 Forms).  
21  
22

### 23 24 25 26 27 *Socio-demographic and biophysical measures*

28  
29 Socio-demographic and biophysical data of respondents were recorded by using a predesigned  
30 questionnaire. The collected information was regarding their age, gender, body mass index  
31 (BMI), marital status, education level, occupation, economic status, residence, living status  
32 (with or without family), and smoking habit.  
33  
34  
35

### 36 37 38 *UCLA loneliness scale (ULS-8)*

39  
40 The ULS-8 is a short form of the UCLA loneliness scale containing 8 questions.<sup>21</sup> Each  
41 question has a 4-level score according to the chosen answer as 1 (never), 2 (rarely), 3  
42 (sometimes), 4 (often). Two questions are positively worded, reverse marking from others (“I  
43 am an outgoing person,” and “I can find companionship when I want it”). The total score ranges  
44 from 8 to 32 where a higher score suggested a higher degree of loneliness. We segmented the  
45 total score in 4 groups i.e. 8 to 16 suggested no loneliness, 17-21 depicts mild loneliness, 22-  
46 16 indicates moderate and finally, 27 or above is marked as severe loneliness.  
47  
48  
49  
50

### 51 52 53 *Patient health questionnaire-9 (PHQ-9)*

54  
55 The self-assessment questionnaire PHQ-9 has 9 different questions for the evaluation of the  
56 depression status of respondents for the last 2 weeks. The total score ranges from 0 to 27 points  
57 where each question is scored 0-3 points based on the answers (0, not at all; 1, several days; 2,  
58 half of the days; 3, nearly every day). Below 10 points indicates no depressive symptoms where  
59  
60

1  
2  
3 10 or higher indicates the presence of depressive symptoms.<sup>22</sup> To determine the gravity of  
4 depression, we divided the total score into four levels. Total scores <10, 10-15, 16-21, and 22-  
5 27 for no depression, mild depression, moderate depression, and severe depression,  
6 respectively.  
7  
8  
9

### 10 *Generalized anxiety disorder 7-item (GAD-7) scale*

11  
12  
13 The GAD-7 is a one-dimensional instrument consisting of 7 questions indicating how often,  
14 over the past 2 weeks, the respondents have been bothered by each of the seven key symptoms  
15 of GAD. Total score ranges from 0 to 21 where each question has 4 different scores according  
16 to responses like 0 (not at all), 1 (several days), 2 (more than half the days), and 3 (nearly every  
17 day). A higher score indicates a high degree of anxiety, divided into 4 different segments where  
18 <5, 5-9, 10-14, and 15-21 indicate no, mild, moderate, and severe anxiety, respectively.<sup>23</sup>  
19  
20  
21  
22  
23

### 24 *Pittsburgh sleep quality index (PSQI)*

25  
26 The PSQI questionnaire was used to determine the sleep quality of respondents over one month.  
27 This questionnaire consists of 19 specific questions in 7 different domains: (1) sleep quality (1  
28 question), (2) sleep latency (2 questions), (3) sleep duration (1 question), (4) sleep efficiency  
29 (3 questions), (5) sleep disturbance (9 questions), (6) sleep medication (1 question), and (7)  
30 daily dysfunction (2 questions). Each domain score ranging from 0 to 3 and the total final score  
31 of seven domains is 0 to 21. The higher total scores represent the poor sleep quality,  
32 determining the severity. A total PSQI score below 5 shows no sleep problem and greater than  
33 5 indicates poor sleep. Moreover, respondents having PSQI score greater than 10 are  
34 considered as bad sleepers.<sup>24</sup>  
35  
36  
37  
38  
39  
40  
41

### 42 *Statistical analysis*

43  
44 Microsoft Excel 2016 and statistical packages for social sciences (IBM SPSS, version 25.0)  
45 were used to perform statistical analysis. The data editing, sorting, coding, classification, and  
46 tabulation were performed by Microsoft Excel. Then the excel file was imported into IBM  
47 SPSS software. The descriptive statistics were applied to analyze the characteristics of the  
48 respondents. A chi-square test was performed to observe the differences in loneliness status  
49 (yes or no) with or without depression, anxiety, or sleep problem among the respondents. The  
50 correlations between risk factors and psychometric measures (loneliness, depression, anxiety,  
51 and sleep quality) were assessed with binary logistic regression analysis with a 95% confidence  
52 interval. Statistically significant results were considered at  $p < 0.05$ .  
53  
54  
55  
56  
57  
58  
59  
60

### Patient and public involvement

Patients and the public were not involved in this study.

## RESULTS

The descriptive statistics for all variables of the respondents are presented in [Table 1](#) and [Table 2](#).

Table 1: Distribution of variables and their association with loneliness and depression among the respondents.

	Total N = 672		Loneliness N = 478		$\chi^2$	df	p- value	Depression N = 256		$\chi^2$	df	p- value
			Yes					Yes				
	n	%	n	%				n	%			
<b>Age</b>												
15 to 30 years	344	51	236	69	2.19	1	0.139	119	35	3.67	1	0.056
Above 30 years	328	49	242	74				137	42			
<b>Gender</b>												
Male	381	57	278	73	1.44	1	0.230	117	31	20.35	1	<0.001
Female	291	43	200	69				139	48			
<b>BMI (kg/m<sup>2</sup>)</b>												
Below 18.5	32	5	24	75	0.25	2	0.880	18	56	4.74	2	0.093
18.5–25	435	65	308	71				163	37			
Above 25	205	31	146	71				75	37			
<b>Marital status</b>												
Unmarried	289	43	210	73	0.58	1	0.446	107	37	0.25	1	0.619
Married	383	57	268	70				149	39			
<b>Education</b>												
Illiterate	26	4	21	81	7.53	4	0.110	15	58	6.00	4	0.199
Primary	85	13	59	69				35	41			
Secondary	61	9	48	79				23	38			
Higher Secondary	410	61	279	68				146	36			
Graduate/above	90	13	71	79				37	41			
<b>Occupation</b>												
Service	219	33	153	70	3.75	3	0.290	66	30	12.45	3	0.006
Business	28	4	20	71				7	25			
Student	208	31	158	76				88	42			
Unemployed	217	32	147	68				95	44			
<b>Economic status</b>												
Low	245	36	179	73	0.75	2	0.688	106	43	6.11	2	0.047
Medium	310	46	218	70				103	33			
High	117	17	81	69				47	40			
<b>Residence</b>												
Urban	255	38	184	72	0.21	1	0.646	109	43	3.79	1	0.052
Rural	417	62	194	47				147	35			
<b>Living status</b>												

With family	430	64	291	68	11.62	3	<b>0.009</b>	179	42	9.86	3	<b>0.020</b>
Without family	242	36	187	77				77	32			
<b>Smoking habit</b>												
Smoker	177	26	135	76	3.09	1	0.079	50	28	9.88	1	<b>0.002</b>
Non-smoker	495	74	343	69				206	42			
<b>Loneliness</b>												
Yes	478	71	478	10	-	-	-	228	48	64.75	1	<b>&lt;0.001</b>
No	194	29	0	0				28	14			
<b>Depression</b>												
Yes	256	38	228	89	64.75	1	<b>&lt;0.001</b>	256	10	-	-	-
No	416	62	250	60				0	0			

Table 2: Distribution of variables and their association with generalized anxiety and sleep problems among the respondents.

	Generalized anxiety N= 429					Sleep problem N = 494				
	Yes		$\chi^2$	df	p-value	Yes		$\chi^2$	df	p-value
	n	%				n	%			
<b>Age</b>										
15 to 30 years	213	62	1.12	1	0.289	242	70	3.62	1	0.051
Above 30 years	216	66				252	77			
<b>Gender</b>										
Male	223	59	10.74	1	<b>0.001</b>	271	71	2.57	1	0.109
Female	206	71				233	80			
<b>BMI (kg/m<sup>2</sup>)</b>										
Below 18.5	28	88	8.16	2	<b>0.017</b>	26	81	1.08	2	0.584
18.5–25	272	63				317	73			
Above 25	129	63				151	74			
<b>Marital status</b>										
Unmarried	187	65	0.16	1	0.685	207	72	0.93	1	0.336
Married	242	63				287	75			
<b>Education</b>										
Illiterate	18	69	7.39	4	0.117	19	73	0.71	4	0.849
Primary	61	72				65	76			
Secondary	44	72				44	72			
Higher Secondary	246	60				302	74			
Graduate/above	60	67				64	71			
<b>Occupation</b>										
Service	125	57	9.28	3	<b>0.026</b>	158	72	0.56	3	0.906
Business	17	61				20	71			
Student	148	71				153	74			
Unemployed	139	64				163	75			
<b>Economic status</b>										
Low	163	67	1.39	2	0.499	177	72	2.60	2	0.273
Medium	195	63				224	72			
High	71	61				93	79			

<b>Residence</b>										
Urban	164	64	0.04	1	0.841	192	75	0.67	1	0.413
Rural	265	64				302	72			
<b>Living status</b>										
With family	274	64	1.56	3	0.665	323	75	4.82	3	0.185
Without family	155	64				171	71			
<b>Smoking habit</b>										
Smoker	105	59	2.12	1	0.145	129	73	0.05	1	0.825
Non-smoker	324	65				365	74			
<b>Generalized anxiety</b>										
Yes	429	100	-	-	-	366	85	84.86	1	<b>&lt;0.001</b>
No	0	0				128	53			
<b>Sleep problem</b>										
Yes	366	75	84.88	1	<b>&lt;0.001</b>	494	100	-	-	-
No	63	35				0	0			

BMI: Body mass index; df: Degrees of freedom; N: Number.

The total respondents were 672, among them, males and females were 57% and 43%, respectively. About half of the total respondents were above 30 years of age and half were below 30 years. Sixty-five percent of total respondents had normal BMI and 61% of respondents completed their higher secondary education. The majority of the respondents were married (57%), non-smoker (74%), and medium economic class (46%). We observed two-third of the total respondents are living with family (64%) in a rural area (62%).

The estimations of loneliness, depression, anxiety, and sleep problems were 71%, 38%, 64%, and 74%, respectively (Figure 1). The proportions of respondents having loneliness were higher in (i) individuals living without family members vs. with family members (77% vs. 68%,  $p=0.009$ ), (ii) individuals with vs. without considerable depression (89% vs. 60%,  $p<0.001$ ), anxiety (85% vs. 46%,  $p<0.001$ ), and sleep problem (79% vs. 49%,  $p<0.001$ ), respectively. The sizes of respondents with depression were higher in (i) females vs. males (48% vs. 31%,  $p < 0.001$ ), (ii) unemployed vs. service (44% vs. 30%,  $p=0.006$ ), (iii) non-smoker vs. smoker (42% vs. 28%,  $p=0.002$ ), (iv) individuals with economic status lower vs. medium (43% vs. 33%,  $p=0.047$ ), (v) respondents living with family members vs. without family members (42% vs. 33%,  $p = 0.020$ ), (vi) individuals with vs. without considerable loneliness (48% vs. 14%,  $p<0.001$ ), anxiety (57% vs. 5%,  $p<0.001$ ), and sleep problem (45% vs. 19%,  $p<0.001$ ), respectively. The extents of respondents with anxiety were higher in (i) females vs. males (71% vs. 69%,  $p=0.001$ ), (ii) students vs. service (71% vs. 57%,  $p=0.026$ ), (iii) individuals with BMI below 18.5 vs. above 25 kg/m<sup>2</sup> (88% vs. 63%,  $p=0.017$ ), (iv) individuals with vs. without considerable loneliness (77% vs. 32%,  $p<0.001$ ), depression (95% vs. 44%,  $p<0.001$ ), and sleep problem (75% vs. 35%,  $p<0.001$ ), respectively. And, finally the

proportions of respondents with sleep problem were higher in individuals with vs. without considerable loneliness (82% vs. 53%,  $p < 0.001$ ), depression (87% vs. 65%,  $p < 0.001$ ), and anxiety (85% vs. 53%,  $p < 0.001$ ), respectively.

The binary logistic regression analysis was performed to measure the correlations between dependent and independent variables and demonstrated in [Table 3](#) and [Table 4](#).

Table 3: Regression analysis of variables by mental health problems (Loneliness and Depression) among respondents.

	Loneliness N = 478			Depression N = 256		
	OR	95% CI	P-value	OR	95% CI	p-value
<b>Age</b>						
15 to 30 years	0.606	0.359-1.023	0.061	0.653	0.386-1.105	0.112
Above 30 years	1			1		
<b>Gender</b>						
Male	0.666	0.411-1.080	0.099	1.624	1.000-2.637	<b>0.050</b>
Female	1			1		
<b>BMI (kg/m<sup>2</sup>)</b>						
Below 18.5	1.574	0.557-4.443	0.392	0.889	0.346-2.282	0.806
18.5-25	1.244	0.399-3.877	0.706	0.832	0.287-2.410	0.735
Above 25	1			1		
<b>Marital status</b>						
Unmarried	0.902	0.508-1.600	0.723	0.845	0.463-1.542	0.584
Married	1			1		
<b>Education</b>						
Illiterate	0.498	0.211-1.172	0.111	1.076	0.476-2.430	0.861
Primary	1.365	0.491-3.794	0.551	1.545	0.620-3.851	0.351
Secondary	1.123	0.293-4.309	0.866	1.813	0.507-6.485	0.360
Higher Secondary	0.518	0.204-1.314	0.166	1.006	0.430-2.351	0.989
Graduate/above	1			1		
<b>Occupation</b>						
Service	0.889	0.292-2.711	0.836	0.315	0.099-1.009	0.052
Business	1.476	0.895-2.434	0.127	1.760	1.037-2.994	0.036
Student	0.628	0.264-1.962	0.253	0.752	0.363-1.052	0.093
Unemployed	1			1		
<b>Economic status</b>						
Low	1.111	0.636-1.941	0.712	1.478	0.832-2.624	0.183
Medium	1.133	0.725-1.771	0.583	1.618	1.038-2.252	<b>0.033</b>
High	1			1		
<b>Residence</b>						
Urban	1.060	0.697-1.611	0.785	0.655	0.434-0.988	<b>0.044</b>
Rural	1			1		
<b>Living status</b>						
With family	0.457	0.278-0.751	<b>0.002</b>	1.797	1.126-2.867	<b>0.014</b>
Without family	1			1		
<b>Smoking habit</b>						

Smoker	0.637	0.376-1.078	0.093	1.441	0.849-2.447	0.176
Non-smoker	1			1		
<b>Loneliness</b>						
Yes	-	-	-	2.890	1.669-5.000	<0.001
No	-			1		
<b>Depression</b>						
Yes	2.958	1.733-5.050	<0.001	-	-	-
No	1			-		
<b>Generalized anxiety</b>						
Yes	3.952	2.506-6.211	<0.001	18.518	9.523-35.714	<0.001
No	1			1		
<b>Sleep problem</b>						
Yes	2.638	1.718-4.048	<0.001	1.385	0.814-2.352	0.229
No	1			1		

Table 4: Regression analysis of variables by mental health problems (Generalized anxiety and Sleep problem) among respondents.

	Generalized anxiety N= 429			Sleep disorder N = 494		
	OR	95% CI	p-value	OR	95% CI	p-value
<b>Age</b>						
15 to 30 years	0.857	0.490-1.501	0.590	0.834	0.499-1.394	0.489
Above 30 years	1			1		
<b>Gender</b>						
Male	1.453	0.869-2.431	0.154	1.036	0.648-1.658	0.882
Female	1			1		
<b>BMI (kg/m<sup>2</sup>)</b>						
Below 18.5	0.270	0.072-1.011	0.052	0.818	0.289-2.319	0.706
18.5–25	0.347	0.085-1.411	0.139	0.877	0.281-2.737	0.821
Above 25	1			1		
<b>Marital status</b>						
Unmarried	1.097	0.599-2.011	0.763	1.009	0.578-1.764	0.974
Married	1			1		
<b>Education</b>						
Illiterate	0.511	0.215-1.211	0.127	1.346	0.617-2.935	0.455
Primary	0.566	0.210-1.525	0.260	0.855	0.350-2.090	0.732
Secondary	0.244	0.059-1.006	0.051	0.829	0.249-2.763	0.761
Higher Secondary	0.788	0.311-2.000	0.616	1.396	0.589-3.310	0.449
Graduate/above	1			1		
<b>Occupation</b>						
Service	0.855	0.296-2.481	0.774	0.894	0.318-2.515	0.831
Business	1.300	0.754-2.237	0.344	1.161	0.702-1.923	0.559
Student	0.548	0.362-0.952	0.452	0.524	0.241-1.183	0.263
Unemployed	1					
<b>Economic status</b>						
Low	0.562	0.312-1.011	0.054	1.588	0.893-2.824	0.115

Medium	0.981	0.613-1.572	0.938	0.852	0.558-1.300	0.458
High	1			1		
<b>Residence</b>						
Urban	1.146	0.737-1.781	0.545	0.929	0.618-1.397	0.724
Rural	1			1		
<b>Living status</b>						
With family	0.842	0.511-1.388	0.501	1.344	0.844-2.140	0.213
Without family	1			1		
<b>Smoking habit</b>						
Smoker	1.032	0.611-1.743	0.906	0.925	0.567-1.509	0.755
Non-smoker	1			1		
<b>Loneliness</b>						
Yes	3.952	2.500-6.211	<0.001	2.525	1.647-3.875	<0.001
No	1			1		
<b>Depression</b>						
Yes	18.518	9.615-35.714	<0.001	1.344	0.800-2.257	0.263
No	1			1		
<b>Generalized anxiety</b>						
Yes	-	-	-	3.367	2.127-3.519	<0.001
No	-	-	-	1		
<b>Sleep problem</b>						
Yes	3.558	2.237-5.649	<0.001	-	-	-
No	1			-		

BMI: Body mass index; CI: Confidence interval; N: Number; OR: Odds ratio.

Respondents living with family members were 0.46 times less likely than the respondents living away from family members to have loneliness (OR = 0.46; 95% CI = 0.28-0.75,  $p=0.002$ ). Respondents having business occupations were 1.76 times more likely than the unemployed group to have depression (OR = 1.76; 95% CI = 1.04–2.99,  $p=0.036$ ). Respondents from the medium economic class were 1.62 times more likely than those who were from high economic impression to have depression (OR = 1.62; 95% CI = 1.04–2.25,  $p=0.033$ ). Urban respondents were 0.66 times less likely to have depression than rural respondents (OR = 0.66; 95% CI = 0.43-0.99,  $p=0.044$ ). Respondents who are living with family members were 1.80 times more likely to have depression than those of the respondents who are living away from the family members (OR = 1.80; 95% CI = 1.13-2.87,  $p=0.014$ ). The probabilities of having loneliness were 2.96 times higher in respondents who already have depression (OR = 2.96; 95% CI = 1.73-5.05,  $p<0.001$ ), 3.95 times higher in respondents with anxiety (OR = 3.95; 95% CI = 2.51-6.21,  $p<0.001$ ), and 2.64 times higher in respondents suffering from sleep problem (OR = 2.64; 95% CI = 1.72-4.05,  $p<0.001$ ), respectively. The chances of having depression were 18.51 times higher in respondents with anxiety (OR = 18.51; 95% CI = 9.52-35.71,  $p<0.001$ ), and 1.36 times higher in respondents suffering from sleep problem (OR = 1.36; 95% CI = 0.81-2.35,  $p<0.001$ ), respectively. The likelihoods of



1  
2  
3 having anxiety were 3.60 times higher in respondents suffering from sleep problems (OR =  
4 3.60; 95% CI = 2.24-5.65,  $p < 0.001$ ).  
5  
6

## 7 **DISCUSSION**

8  
9  
10 The present prospective study demonstrates the impact of pandemic COVID-19 on the mental  
11 health status among the general population of Bangladesh. The spreading of COVID-19 is  
12 spiraling out of control worldwide; the general population around the world is dreadfully  
13 frazzled currently.<sup>25</sup> The psychological reactions of the population toward any outbreak of  
14 infectious diseases play an important role in controlling both the spreading of the disease and  
15 the occurrence of emotional disturbances and in the regulation of social disorder during and  
16 after the pandemic.<sup>26</sup> The general population is always under stressful conditions because of  
17 the unpredictability of the situations, when to control the spreading of the disease, and the  
18 seriousness of the risk factors associated with COVID-19.<sup>27</sup> Experiences from past similar  
19 epidemics and pandemics, serious concerns such as fear of death arise among the patients, and  
20 feelings of loneliness, anger, depression, sleep disturbance can develop among people who are  
21 quarantined or self-isolated.<sup>9</sup> It can be inferred that the general population who are in lockdown  
22 might develop several mental health problems.  
23  
24  
25  
26  
27  
28  
29  
30  
31

32  
33 The government-imposed lockdown situation or stay home orders has interrupted the normal  
34 life of general people that ultimately impacted the individual's mental health.<sup>28 29</sup> General  
35 people are living a monotonous, unusual, helpless life in their locked home, blaming others to  
36 be responsible that slowly developing mental breakdown.<sup>30</sup> Patients of COVID-19 or suspected  
37 isolated people are more vulnerable to experience trauma due to fear of death. Suspected  
38 patients who are staying in quarantine can feel loneliness, depression, anxiety, sleep problems,  
39 bored, and panic that may lead to emotional disturbance, self-harm, substance abuse, and  
40 suicidal thoughts.<sup>31 32</sup> A rapid review found that stress factor indefinite lockdown conditions,  
41 fear of infection, frustration, monotonous life, insufficient supplies, inadequate information,  
42 financial loss, and fear of an uncertain future have led to developing long-lasting posttraumatic  
43 stress symptoms, loneliness, anxiety, confusion, anger, depression among the general  
44 population.<sup>33</sup>  
45  
46  
47  
48  
49  
50  
51  
52  
53

54 The present study found that about three in every four people in Bangladesh are suffering from  
55 loneliness and sleep problems at any level of gravity during the lockdown period (Figure 1).  
56 People with considerable loneliness are prone to have sleep problems than without loneliness.  
57 On the other hand, people living without family during lockdown are more prone to develop  
58  
59  
60

1  
2  
3 loneliness than people living with family. Following the current findings, one study conducted  
4 in the United Kingdom and observed 36% of the respondents feel lonely during the COVID-  
5 19 pandemic which is quite higher than any normal times.<sup>34</sup> Another two studies also reported  
6 higher loneliness scores among general people during COVID-19 than past normal times.<sup>35 36</sup>  
7  
8 We also found that depression is higher in females than males like previous findings.<sup>37-40</sup> The  
9  
10 prevalence of depression and generalized anxiety among the Bangladeshi population during  
11 the early lockdown period was 38% and 64%, respectively. The increased rate of depression  
12 was observed in several recent studies in Spain, China, and Hong Kong due to the COVID-19  
13 pandemic.<sup>41-43</sup> Also, the high prevalence anxiety disorder was observed among respondents  
14 from many countries in the COVID-19 pandemic than previous times.<sup>3 37 38</sup> The extent of  
15 anxiety among Bangladeshi students found about 71% in the present study. Final year students  
16 of different education levels are more anxious about their uncertain exams and the job market.  
17 The present study shows that respondents having loneliness, depression, or anxiety are more  
18 likely to have sleep problems than healthy individuals. Similar findings were reported in many  
19 recent studies. During the COVID-19 pandemic, 29.2% for insomnia, 27.9% for depression,  
20 and 31.6% anxiety was observed in China.<sup>44</sup> During COVID-19 lockdown, 42.2% sleep  
21 disturbances, 17.4% moderate or severe insomnia, 24.7%, and 23.2% depression and anxiety  
22 symptom was observed in Italy.<sup>45</sup>  
23  
24

25  
26 To prevent the quick spread of COVID-19 infection, the Bangladesh government postponed  
27 all educational institutions since March 18, 2020, and imposed a countrywide lockdown to limit  
28 public movement and to inspire staying at home since March 26, 2020.<sup>14</sup> The nationwide  
29 lockdown, restriction on movement or stay home orders have a profound impact on the  
30 economy in the markets, offices, business organizations, and transport systems.<sup>46 47</sup> Most of the  
31 Bangladeshi population mainly depends on a regular income but they are uncertain about  
32 returning to the workplace due to indefinite lockdown, these are producing a confounding  
33 impact on their mental health status.<sup>30</sup> Several suicide cases are reported in Bangladesh due to  
34 getting infected and lost economic opportunities during the lockdown period informing  
35 socioeconomic challenges of marginalized people that reflect the mental health status of  
36 Bangladeshi wage-earners.<sup>48 49</sup> An Indian case study suggested that COVID-19 may  
37 significantly impact the mental health status and influence suicidal ideation, suicide attempts  
38 among the affected people; other comorbid diseases may aggravate the situation.<sup>50</sup> We found  
39 that 44% of unemployed and 30% of service-holders were suffering from depression. About  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 43% of people belonging to the lower economic class are facing depression during the  
4 lockdown period.  
5

6  
7 National Institute of Mental Health (NIMH) of Bangladesh have announced some  
8 recommendations such as psychotropics, avoid COVID-19 news multiple times in a day or  
9 news scrolling, one or two authenticate source of information, less social media, simple  
10 relaxation technique like breathing exercise, etc. regarding the management of mental health  
11 during COVID-19 pandemic.<sup>51</sup> In agreement with the above suggestions, several studies also  
12 recommended online response, counseling, social support, training on mental health for the  
13 patients, health care professionals, public service holders, youth, students, elderly populations  
14 to manage mental health problems during and after the pandemic.<sup>32 52 53</sup> Worldwide, the  
15 COVID-19 pandemic unveiled the actual setup of unpreparedness of the healthcare systems  
16 and the scarcity of resources (PPE, testing kits, etc.) to combat the situation.<sup>54</sup>  
17  
18  
19  
20  
21  
22  
23  
24

### 25 **Strengths and limitations of this study**

26  
27 The present study has a few potential limitations, which should be considered. First, compared  
28 to face-face interviews, online self-reporting surveys have multiple biases and are not  
29 representative for those who have no internet facilities. Secondly, the cross-sectional nature of  
30 the study is not able to measure the impact of these mental illnesses over time. Additionally,  
31 the present study didn't consider the altered life-style of the respondents due to the COVID-19  
32 pandemic. Besides the limitations, the present study has some strong advantages. This is the  
33 first-ever study carried out in Bangladesh intending to identify four important mental health  
34 problems (loneliness, depression, anxiety, and sleep problems) during this pandemic time. This  
35 study will provide an idea about the potential mental health situation among the general  
36 Bangladeshi population during the COVID-19 pandemic.  
37  
38  
39  
40  
41  
42  
43  
44

### 45 **Future research**

46  
47 This study highlights the need for mental health assessment and proper management of these  
48 issues during COVID-19 pandemic. Further research among the healthcare professionals to  
49 explore their actual mental health status in this pandemic situation.  
50  
51  
52

### 53 **CONCLUSION**

54  
55 In summary, our finding reflects the gravity of mental health problems during the COVID-19  
56 pandemic where a large portion of general people in Bangladesh affected mentally with  
57 different levels of severity. Loneliness, depression, generalized anxiety, and sleep problems  
58  
59  
60

1  
2  
3 were linked with specific factors and we suggest intensive mental healthcare services for the  
4 Bangladeshi people would support them to reduce these mental health problems during and  
5 after the COVID-19 pandemic. Therefore, the government, non-government, and community  
6 activities should be integrated to ensure individual and collective physical and mental health.  
7  
8 In addition to medical care, social security, economic efficiency, the resilience of affected  
9 groups, strengthening healthcare facilities should be the top priority to grow confidence among  
10 the general population of Bangladesh after the COVID-19 pandemic.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

## Declarations

## Acknowledgments

All the authors are thankful to the participants for their cooperation to conduct this study.

## Author contributions

RD conceived the idea, designed the study, collected the data, and prepared the initial draft of the manuscript. MRH and SD contributed to the development of the study design, collected the data, and edited the manuscript. MRI conceived and designed the study, analyzed and interpreted the data, reviewed the manuscript for important intellectual content, and supervised the whole work. All the authors approved the final version of the manuscript for this submission.

## Funding

The study was self-funded.

## Competing interests

The authors do not have any competing interests to declare.

## Patient consent for publication

Not applicable.

## Data availability statement

Data supporting the present study findings are within this article. All the relevant data and information can be obtained from the corresponding author upon reasonable request.

## Table Legends

**Table 1** Distribution of variables and their association with loneliness and depression among the respondents.

BMI: Body mass index; df: Degrees of freedom; N: Number.

**Table 2** Distribution of variables and their association with generalized anxiety and sleep problems among the respondents.

BMI: Body mass index; df: Degrees of freedom; N: Number.

**Table 3** Regression analysis of variables by loneliness and depression among the study population.

BMI: Body mass index; CI: Confidence interval; N: Number; OR: Odds ratio.

**Table 4** Regression analysis of variables by generalized anxiety and sleep problems among the study population.

BMI: Body mass index; CI: Confidence interval; N: Number; OR: Odds ratio.

## Figure Legends

**Figure 1:** Mental health problems among the respondents based on their distribution and gravity.

## REFERENCES

1. Talevi D, Socci V, Carai M, et al. Mental health outcomes of the COVID-19 pandemic. *Rivista di Psichiatria*. 2020;55(3):137-144.
2. Worldometer. Coronavirus cases, 2020. Available: [www.worldometers.info/coronavirus/](http://www.worldometers.info/coronavirus/) [accessed October 9, 2020].
3. Chen Y, Zhou H, Zhou Y, Zhou F. Prevalence of self-reported depression and anxiety among pediatric medical staff members during the COVID-19 outbreak in Guiyang, China. *Psychiatry Res*. 2020;288:113005.
4. Duan L, Zhu G. Psychological interventions for people affected by the COVID-19 epidemic. *Lancet Psychiatry*. 2020;7(4):300-302.
5. Madabhavi I, Sarkar M, Kadakol N. COVID-19: a review. *Monaldi Arch Chest Dis* 2020;90(2).
6. Zhang Y, Ma ZF. Impact of the COVID-19 Pandemic on Mental Health and Quality of Life among Local Residents in Liaoning Province, China: A Cross-Sectional Study. *Int J Environ Res Public Health*. 2020;17(7):2381.
7. Maunder RG, Lancee WJ, Balderson KE, et al. Long-term psychological and occupational effects of providing hospital healthcare during SARS outbreak. *Emerg Infect Dis*. 2006;12(12):1924-1932.
8. North CS, Pfefferbaum B, Vythilingam M, et al. Exposure to bioterrorism and mental health response among staff on Capitol Hill. *Biosecur Bioterror*. 2009;7(4):379-388.
9. Xiang YT, Yang Y, Li W, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry*. 2020;7(3):228-229.
10. Central intelligence agency. The world factbook, 2020. Available: [https://www.cia.gov/library/publications/resources/the-world-factbook/geos/print\\_bg.html](https://www.cia.gov/library/publications/resources/the-world-factbook/geos/print_bg.html) [Accessed 01 September 2020].
11. Streatfield PK, Karar ZA. Population challenges for Bangladesh in the coming decades. *J Health Popul Nutr*. 2008;26(3):261-272.
12. Paul R. Bangladesh Confirms its First Three Cases of Coronavirus, 2020. Available: <https://www.reuters.com/article/us-healthcoronavirus-bangladesh-idUSKBN20V0FS> [Accessed 14 April 2020].
13. Directorate General of Health Services. The latest status in Bangladesh: Covid-19, 2020. Available:

- 1  
2  
3 [https://corona.gov.bd/?gclid=EAIaIQobChMIiL5udzH6wIVU3ZgCh0VYAxHEAAYASAAEgKEAPD\\_BwE](https://corona.gov.bd/?gclid=EAIaIQobChMIiL5udzH6wIVU3ZgCh0VYAxHEAAYASAAEgKEAPD_BwE) [Accessed 08 October 2020].
- 4  
5  
6  
7 14. The Daily Star. Man left out of relief net dies by suicide in Chattogram, 2020. Available:  
8 [https://www.thedailystar.net/coronavirus-deadly-new-threat/news/man-left-out-relief-net-](https://www.thedailystar.net/coronavirus-deadly-new-threat/news/man-left-out-relief-net-dies-suicide-chattogram-1893409)  
9 [dies-suicide-chattogram-1893409](https://www.thedailystar.net/coronavirus-deadly-new-threat/news/man-left-out-relief-net-dies-suicide-chattogram-1893409) [Accessed 16 April 2020].
- 10  
11  
12 15. The Kaler Kantho. Day labourer commits suicide in Naldanga (In Bangla), 2020. Available:  
13 <https://www.kalerkantho.com/home/printnews/898402/2020-04-13> [Accessed 13 April  
14 2020].
- 15  
16  
17 16. The Jugantor. The young woman committed suicide after losing her job in Corona left  
18 posting on Facebook (In Bangla), 2020. Retrieved July 11, 2020, Available:  
19 <https://rb.gy/pmssav> [Accessed 11 July 2020].
- 20  
21  
22 17. Banna MHA, Sayeed A, Kundu S, et al. The impact of the COVID-19 pandemic on the  
23 mental health of the adult population in Bangladesh: a nationwide cross-sectional study  
24 [published online ahead of print, 2020 Aug 2]. *Int J Environ Health Res.* 2020;1-12.
- 25  
26  
27 18. Islam MS, Ferdous MZ, Potenza MN. Panic and generalized anxiety during the COVID-  
28 19 pandemic among Bangladeshi people: An online pilot survey early in the outbreak  
29 [published online ahead of print, 2020 Jul 15]. *J Affect Disord.* 2020;276:30-37.
- 30  
31  
32 19. Khan AH, Sultana MS, Hossain S, Hasan MT, Ahmed HU, Sikder MT. The impact of  
33 COVID-19 pandemic on mental health & wellbeing among home-quarantined Bangladeshi  
34 students: A cross-sectional pilot study [published online ahead of print, 2020 Aug 7]. *J*  
35 *Affect Disord.* 2020;277:121-128.
- 36  
37  
38 20. Yeasmin S, Banik R, Hossain S, et al. Impact of COVID-19 pandemic on the mental health  
39 of children in Bangladesh: A cross-sectional study. *Child Youth Serv Rev.*  
40 2020;117:105277.
- 41  
42  
43 21. Hays RD, DiMatteo MR. A short-form measure of loneliness. *J Pers Assess.*  
44 1987;51(1):69-81.
- 45  
46  
47 22. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity  
48 measure. *J Gen Intern Med.* 2001;16(9):606-613.
- 49  
50  
51 23. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized  
52 anxiety disorder: the GAD-7. *Arch Intern Med.* 2006;166(10):1092-1097.
- 53  
54  
55 24. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta psychiatrica*  
56 *scandinavica.* 1983;67(6):361-370.
- 57  
58  
59 25. Dubey S, Biswas P, Ghosh R, et al. Psychosocial impact of COVID-19 [published online  
60 ahead of print, 2020 May 27]. *Diabetes Metab Syndr.* 2020;14(5):779-788.



- 1
- 2
- 3 26. Taylor S. The psychology of pandemics: Preparing for the next global outbreak of
- 4 infectious disease. Cambridge Scholars Publishing; 2019.
- 5
- 6 27. Bao Y, Sun Y, Meng S, Shi J, Lu L. 2019-nCoV epidemic: address mental health care to
- 7 empower society. *Lancet*. 2020;395(10224):e37-e38.
- 8
- 9 28. Nicola M, Alsaifi Z, Sohrabi C, et al. The socio-economic implications of the coronavirus
- 10 pandemic (COVID-19): A review. *Int J Surg*. 2020;78:185-193.
- 11
- 12 29. Holmes EA, O'Connor RC, Perry VH, et al. Multidisciplinary research priorities for the
- 13 COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry*.
- 14 2020;7(6):547-560.
- 15
- 16 30. Hossain KA, Roy S, Ullah MM, Kabir R, Arafat SY. COVID-19 and Mental Health
- 17 Challenges in Bangladesh. *Advances in Medical, Dental and Health Sciences*. 2020;2(3)31-
- 18 3.
- 19
- 20 31. Koh D. Occupational risks for COVID-19 infection. *Occup Med (Lond)*. 2020;70(1):3-5.
- 21
- 22 32. Wang C, Pan R, Wan X, et al. Immediate Psychological Responses and Associated Factors
- 23 during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the
- 24 General Population in China. *Int J Environ Res Public Health*. 2020;17(5):1729.
- 25
- 26 33. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how
- 27 to reduce it: rapid review of the evidence. *Lancet*. 2020;395(10227):912-920.
- 28
- 29 34. Li LZ, Wang S. Prevalence and predictors of general psychiatric disorders and loneliness
- 30 during COVID-19 in the United Kingdom [published online ahead of print, 2020 Jun 30].
- 31 *Psychiatry Res*. 2020;291:113267.
- 32
- 33 35. Luchetti M, Lee JH, Aschwanden D, et al. The trajectory of loneliness in response to
- 34 COVID-19 [published online ahead of print, 2020 Jun 22]. *Am Psychol*.
- 35 2020;10.1037/amp0000690.
- 36
- 37 36. Killgore WDS, Cloonan SA, Taylor EC, Dailey NS. Loneliness: A signature mental health
- 38 concern in the era of COVID-19. *Psychiatry Res*. 2020;290:113117.
- 39
- 40 37. Zhou SJ, Zhang LG, Wang LL, et al. Prevalence and socio-demographic correlates of
- 41 psychological health problems in Chinese adolescents during the outbreak of COVID-19.
- 42 *Eur Child Adolesc Psychiatry*. 2020;29(6):749-758.
- 43
- 44 38. Islam MR, Ali S, Karmoker JR, et al. Evaluation of serum amino acids and non-enzymatic
- 45 antioxidants in drug-naïve first-episode major depressive disorder. *BMC Psychiatry*.
- 46 2020;20(1):333.
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1  
2  
3 39. Li G, Miao J, Wang H, et al. Psychological impact on women health workers involved in  
4 COVID-19 outbreak in Wuhan: a cross-sectional study. *J Neurol Neurosurg Psychiatry*.  
5 2020;91(8):895-897.  
6
- 7  
8 40. Kang L, Ma S, Chen M, et al. Impact on mental health and perceptions of psychological  
9 care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease  
10 outbreak: A cross-sectional study. *Brain Behav Immun*. 2020;87:11-17.  
11
- 12 41. González-Sanguino C, Ausín B, Castellanos MÁ, et al. Mental health consequences during  
13 the initial stage of the 2020 Coronavirus pandemic (COVID-19) in Spain. *Brain Behav*  
14 *Immun*. 2020;87:172-176.  
15
- 16 42. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality  
17 during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res*.  
18 2020;288:112954.  
19
- 20 43. Choi EPH, Hui BPH, Wan EYF. Depression and Anxiety in Hong Kong during COVID-  
21 19. *Int J Environ Res Public Health*. 2020;17(10):3740.  
22
- 23 44. Shi L, Lu ZA, Que JY, et al. Prevalence of and Risk Factors Associated With Mental Health  
24 Symptoms Among the General Population in China During the Coronavirus Disease 2019  
25 Pandemic. *JAMA Netw Open*. 2020;3(7):e2014053.  
26
- 27 45. Gualano MR, Lo Moro G, Voglino G, Bert F, Siliquini R. Effects of Covid-19 Lockdown  
28 on Mental Health and Sleep Disturbances in Italy. *Int J Environ Res Public Health*.  
29 2020;17(13):4779.  
30
- 31 46. The Bangla Tribune. The patient committed suicide by jumping from the roof as he was  
32 infected with corona (In Bangla), 2020. Available: <https://rb.gy/jfогjx> [Accessed 11 June  
33 2020].  
34
- 35 47. The Somoy News. Patient commits suicide without getting treatment in hospital (In  
36 Bangla), 2020. Available: <https://www.somoynews.tv/pages/details/214750> [Accessed 20  
37 May 2020].  
38
- 39 48. The BBC News. Corona virus: COVID-19 patient committed suicide after escaping from  
40 the hospital (In Bangla), 2020. Available: <https://www.bbc.com/bengali/news-53125017>  
41 [Accessed 21 June 2020].  
42
- 43 49. The Desh Rupantor. Corona-infected businessman commits suicide (In Bangla), 2020.  
44 Available: <https://www.deshrupantor.com/mofossil/2020/07/07/230368> [Accessed 07  
45 July 2020].  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
50. Hossain MM, Tasnim S, Sultana A, McKyer EL, Ma P. COVID-19 and suicide of an army soldier in India: perspectives on psychosocial epidemiology of suicidal behavior. SocArXiv. 2020.
51. National Institute of Mental Health (MIMH). Recommendations from Sub Committee for Mental Health, 2020. Available: [https://mofa.portal.gov.bd/sites/default/files/files/mofa.portal.gov.bd/page/ad1f289c\\_47cf\\_4f6c\\_8dee\\_887957be3176/Guideline%20for%20Mental%20Health%20Issue-COVID-19-%20DGHS.pdf](https://mofa.portal.gov.bd/sites/default/files/files/mofa.portal.gov.bd/page/ad1f289c_47cf_4f6c_8dee_887957be3176/Guideline%20for%20Mental%20Health%20Issue-COVID-19-%20DGHS.pdf) [Accessed 01 September, 2020]
52. Yang Y, Li W, Zhang Q, Zhang L, Cheung T, Xiang YT. Mental health services for older adults in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020;7(4):e19.
53. Armitage R, Nellums LB. The COVID-19 response must be disability inclusive. *Lancet Public Health*. 2020;5(5):e257.
54. Pfefferbaum B, North CS. Mental Health and the Covid-19 Pandemic. *N Engl J Med*. 2020;383(6):510-512.




Figure 1: Mental health problems among the respondents based on their distribution and gravity

**Title: Impact of COVID-19 pandemic on mental health among general Bangladeshi population: A cross-sectional study**

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Section/Paragraph
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	Title
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Abstract
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Introduction Paragraphs 1-3
Objectives	3	State specific objectives, including any prespecified hypotheses	Introduction Paragraphs 4
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	Methods Paragraphs 1
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Methods Paragraphs 1
Participants	6	(a) Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls	Methods Paragraphs 1
		(b) For matched studies, give matching criteria and the number of controls per case	Methods Paragraphs 1
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Methods Paragraphs 1
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Methods Paragraphs 2-3
Bias	9	Describe any efforts to address potential sources of bias	N/A
Study size	10	Explain how the study size was arrived at	Methods Paragraphs 1
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Methods Paragraphs 1
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Methods Paragraphs 8
		(b) Describe any methods used to examine subgroups and interactions	Methods Paragraphs 8
		(c) Explain how missing data were addressed	Methods Paragraphs 4
		(d) If applicable, explain how matching of cases and controls was addressed	N/A
		(e) Describe any sensitivity analyses	N/A
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed	Results Paragraphs 1

		eligible, included in the study, completing follow-up, and analysed	
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Results Paragraphs 1 and Table 1
		(b) Indicate number of participants with missing data for each variable of interest	N/A
Outcome data	15*	Report numbers in each exposure category, or summary measures of exposure	Results Paragraphs 2-4 and Table 2

For peer review only

1			
2	Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included
3			Results Paragraphs 2-4 and Table 2
4			
5			
6			
7			(b) Report category boundaries when continuous variables were categorized
8			N/A
9			
10			(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period
11			N/A
12	Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses
13			N/A
14			
15			
16	<b>Discussion</b>		
17	Key results	18	Summarise key results with reference to study objectives
18			Discussion Paragraph 1
19	Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
20			Discussion Paragraph 5
21			
22			
23	Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
24			Discussion Paragraph 2-4
25			
26			
27	Generalisability	21	Discuss the generalisability (external validity) of the study results
28			Discussion Paragraph 2-4
29	<b>Other information</b>		
30	Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based
31			A small seed fund for students was available.
32			
33			
34			

\*Give information separately for cases and controls.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at <http://www.strobe-statement.org>.

# BMJ Open

## Impact of COVID-19 pandemic on mental health among general Bangladeshi population: A cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2020-045727.R1
Article Type:	Original research
Date Submitted by the Author:	16-Feb-2021
Complete List of Authors:	Das, Rajesh; University of Asia Pacific, Department of Pharmacy Hasan, Md. Rakib; Jahangirnagar University, Department of Pharmacy Daria, Sohel; University of Asia Pacific, Department of Pharmacy Islam, Md. Rabiul; University of Asia Pacific,
<b>Primary Subject Heading</b>:	Public health
Secondary Subject Heading:	Mental health, Health policy, Global health, Epidemiology
Keywords:	COVID-19, Depression & mood disorders < PSYCHIATRY, Anxiety disorders < PSYCHIATRY, Adult psychiatry < PSYCHIATRY, PUBLIC HEALTH

SCHOLARONE™  
Manuscripts





I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1  
2  
3  
4 **Title:** Impact of COVID-19 pandemic on mental health among general  
5 Bangladeshi population: A cross-sectional study  
6  
7  
8  
9  
10  
11

12 **Running Title:** COVID-19 pandemic and mental health issues  
13  
14  
15  
16  
17  
18

19 **Authors:**

20 Rajesh Das<sup>1</sup>, Md. Rakib Hasan<sup>2</sup>, Sohel Daria<sup>1</sup>, Md. Rabiul Islam<sup>1,3\*</sup>  
21  
22  
23  
24  
25  
26

27 **Affiliations:**

28  
29 <sup>1</sup>Department of Pharmacy, University of Asia Pacific, 74/A Green Road, Farmgate, Dhaka-  
30 1215, Bangladesh  
31  
32

33 <sup>2</sup>Department of Pharmacy, Jahangirnagar University, Savar, Dhaka-1335  
34  
35

36 <sup>3</sup>Department of Clinical Pharmacy and Pharmacology, Dhaka University, Dhaka-1000  
37  
38  
39  
40  
41  
42

43 **Word Count: 3,439**

44 *(From the introduction to conclusion excluding abstract, references, declarations, etc.)*  
45  
46  
47

48 **\*Correspondence:**

49 **Md. Rabiul Islam, PhD**

50 Assistant Professor, Department of Pharmacy, University of Asia Pacific

51 74/A Green Road, Farmgate, Dhaka-1215, Bangladesh.

52 Email: robi.ayaan@gmail.com; Cell: +8801916031831

53 ORCID: <https://orcid.org/0000-0003-2820-3144>  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **Title:** Impact of COVID-19 pandemic on mental health among general  
4 Bangladeshi population: A cross-sectional study  
5  
6  
7

8 **ABSTRACT**  
9

10 **Objectives:** The mental health problems are significantly amplified worldwide during the  
11 coronavirus (COVID-19) pandemic. At the early stage of the outbreak, the government of  
12 Bangladesh imposed lockdown and quarantine approaches to prevent virus spread that  
13 impacted people's daily life and health. COVID-19 pandemic also affected people's economic  
14 status, healthcare facilities, and other lifestyle factors in Bangladesh. We aimed to assess the  
15 impact of the COVID-19 pandemic on mental health among the Bangladeshi population.  
16  
17  
18  
19

20  
21 **Methods:** We conducted an online cross-sectional survey among 672 Bangladeshi people aged  
22 between 15 and 65 years all over the country from April 15 to May 10, 2020. After taking  
23 electronic consent, we conducted our survey assessing the socio-demographic profiles and  
24 psychometric measures. We used the UCLA loneliness scale (ULS-8), patient health  
25 questionnaire-9 (PHQ-9), generalized anxiety disorder 7-item (GAD-7) scale, Pittsburgh sleep  
26 quality index (PSQI) for assessment of loneliness, depression, anxiety, and sleep problem,  
27 respectively.  
28  
29  
30  
31  
32

33  
34 **Results:** Estimates of loneliness, depression, anxiety, and sleep problems were 71% (mild:  
35 32%, moderate: 29%, severe: 10%), 38% (mild: 24%, moderate: 11%, severe: 3%), 64% (mild:  
36 30%, moderate: 17%, severe: 17%), and 73% (mild: 50%, moderate: 18%, severe: 5%),  
37 respectively. In Bangladesh, the key factors associated with poor mental health during COVID-  
38 19 were female gender, unemployment, students, obesity, and living without a family. Also,  
39 the present study detected statistically significant inter-relationships among the measured  
40 mental health issues.  
41  
42  
43  
44  
45

46  
47 **Conclusions:** A large portion of respondents reported mental health problems during the  
48 COVID-19 pandemic in Bangladesh. The present study suggests longitudinal assessments of  
49 mental health among Bangladeshi people to know the gravity of this issue during and after the  
50 pandemic. Appropriate supportive programs and interventional approaches would address  
51 mental health problems in Bangladesh during this COVID-19 pandemic.  
52  
53  
54

55 **Keywords:** COVID-19; Mental health; Depression; Anxiety; Loneliness; Pandemics;  
56 Bangladesh  
57  
58  
59  
60

## Article Summary

### Strengths and limitations of this study

- The present study assessed the four major mental health issues among the general Bangladeshi population during the COVID-19 pandemic.
- The present study ensures rapid data collection during public health emergencies and suitability to developing context-specific mental health programs.
- Online self-reporting surveys might have multiple biases and are not representative of those without internet facilities.
- The cross-sectional study is not able to measure the impact of these mental illnesses over time.

## INTRODUCTION

The novel coronavirus strain SARS-CoV-2 causes COVID-19 originated from Wuhan, China. COVID-19 was declared a pandemic on March 11, 2020, due to its uncontrolled spreading worldwide.<sup>1</sup> After spreading, till December 29, 2020, over 79.2 million COVID-19 cases and over 1.7 million deaths were reported worldwide since the start of the pandemic.<sup>2</sup> In December 2020, we observed the highest weekly average of 4.3 million confirmed new cases than any previous time.<sup>3</sup> This devastating condition has not improved yet due to lack of proper treatment and medications though more than a hundred vaccine candidates are in the different developing stages.<sup>4</sup> Most countries-imposed lockdown to limit virus spread that eventually affects people's socioeconomic conditions and mental health regardless of age, gender, profession, etc.<sup>5</sup> Usually, the COVID-19 transmits through breathing droplets or contact with infected individuals. This fear of coronavirus infection impacted the lifestyle, psychology, and relationship status of people.<sup>6</sup> About 52.1% of people felt worried during the COVID-19 pandemic. Among them, 57.8-77.9% wanted mental support from family and friends.<sup>7</sup>

Major epidemic and pandemic outbursts have several negative impacts on individual and collective mental health in society.<sup>5</sup> Previous middle east respiratory syndrome (MERS) and severe acute respiratory syndrome (SARS) severely impacted the local people's mental health.<sup>8</sup> Healthcare providers experienced long-term occupational and psychological effects during the SARS outbreak.<sup>10</sup> In the USA, a study was conducted among Capitol Hill staff workers to respond successfully to disaster-related mental health after the anthrax attacks in 2001. This study reported the prevalence rates of post-traumatic stress disorder and post-anthrax psychiatric disorder among respondents were 55% and 27%, respectively.<sup>11</sup> The mental health of many individuals is potentially affected by COVID-19 in many ways. Family members and friends of COVID-19 patients, their close contacts, isolated or suspected population, healthcare providers, and general people experience extra mental health burdens during the COVID-19 pandemic.<sup>12</sup> Therefore, understanding the impact of the COVID-19 pandemic on an individual's mental health might reduce many current and future mental health issues.

Bangladesh is a densely populated country with a population size of about 164 million people. The population density is five times higher than any other mega country.<sup>13</sup> Several factors that affect mental health are population density, housing, economic status, employment, life experience, disease burden, etc.<sup>14</sup> The economic backbone of the Bangladeshi people squeezed

1  
2  
3 after imposing the lockdown from March 26, 2020. Institute of Epidemiology, Disease Control,  
4 and Research of Bangladesh reported the first COVID-19 cases on March 8, 2020.<sup>15</sup> Between  
5 March 8 and December 27, 2020, there were 509,148 confirmed cases with 7,452 fatalities  
6 reported in Bangladesh. It is the rank 27<sup>th</sup> country contributing 0.64% COVID-19 related  
7 disease burden to the world.<sup>16</sup> At the initial stage, many hospitals were not ready to treat  
8 COVID-19 even the testing facilities were limited. This situation impacted the mental health  
9 of many Bangladeshi people. Few people planned for suicide after failing to cope with these  
10 extra mental burdens.<sup>17-19</sup> The infection has reached every corner of the country. The number  
11 of confirmed cases significantly increased and, mental health-related disorders may increase,  
12 particularly among susceptible people. Therefore, more attention needs to pay to the mental  
13 health burden during and after the COVID-19 pandemic.<sup>20-22</sup> It is equally applicable for other  
14 low- and middle-income countries (LMICs) where resources are limited to tackle any pandemic  
15 situation and its associated mental health issues.<sup>23-25</sup> The present study aimed to assess the  
16 prevalence rates of loneliness, depression, anxiety, and sleep problems among the general  
17 Bangladeshi population during the COVID-19 pandemic. Besides, we expected to find the  
18 associated factors related to mental health problems during the COVID-19 pandemic.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

## 31 **METHODS**

### 32 *Participants and procedure*

33  
34 We carried out a nationwide cross-sectional online survey between 15th April 2020 to 10th  
35 May 2020 using the google survey tool (Google Forms). Here we used the purposive sampling  
36 technique to collect the primary data from the participants. We assumed the confidence  
37 interval, margin of error, and expected prevalence as 95%, 5%, and 30%, respectively.  
38 According to our assumption, the required sample size was 323. Initially, we thought the  
39 response rate might be 20% and invited 1,615 people to participate in the survey. However, the  
40 actual response rate was 46%, we received 736 responses. After screening, we excluded 64  
41 responses due to partial or incomplete information. Finally, we included 672 respondents (381  
42 males, 291 females) aged between 15 to 65 years in the analysis. Before participation in the  
43 survey, the participants reviewed and acknowledged a brief description of the survey, eligibility  
44 requirements, procedures, and e-consent form. We obtained electronic consent from all  
45 participants. All the respondents of this survey were of Bangladeshi ethnicity and living in  
46 Bangladesh at that time. People aged between 15 to 65 years who understood the questions  
47 included in this survey. Exclusion criteria included the history of other psychiatric disorders  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 like delusions, mental retardation, bipolar disorder, schizophrenia, personality disorder, mood-  
4 congruent or incongruent psychotic features, comorbid psychiatric illness, neurological  
5 disease, or patients with clinical evidence of dementia. Additional exclusion criteria were acute  
6 medical conditions, chronic diseases, history of addiction. We did not pay for participation.  
7  
8  
9

### 10 *Estimations*

11  
12 We used two sets of questionnaires for this survey. The first one was a structured questionnaire  
13 designed by the researchers that contained informed consent and socio-demographic  
14 information. The second set was a self-reported structured questionnaire from different  
15 psychometric assessment scales. We prepared both the questionnaire sets in English then  
16 translated them into Bangla. At first, all the questionnaires were translated into Bangla by a  
17 medical graduate and a general person who were native speakers of Bangla and fluent in  
18 English. An independent researcher compiled and addressed the discrepancies to make a single  
19 Bangla forward version. A professional translator having expertise in medical translation and  
20 a medical graduate who was not involved in forward-translation translated this Bangla version  
21 back into English. Again, an independent researcher compiled these back-translated versions  
22 in the same way.<sup>26</sup> We piloted the questionnaire in a randomly selected small group to confirm  
23 clarity and understanding. We circulated the questionnaire in both English and Bengali versions  
24 for the proper understanding of the questions. We sent the link of the designed google form to  
25 participants using email and social networking sites. The educated family members of illiterate  
26 or older respondents helped to collect their responses.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

### 39 *Socio-demographic and biophysical measures*

40  
41 We collected the most relevant socio-demographic information of the respondent. The  
42 collected data were regarding their age, gender, body mass index (BMI), marital status,  
43 education level, occupation, economic status, residence, living status (with or without family),  
44 and smoking habit.  
45  
46  
47  
48

### 49 *UCLA loneliness scale (ULS-8)*

50  
51 The ULS-8 is a short form of the UCLA loneliness scale containing eight questions.<sup>27</sup> Each  
52 question has a 4-level score according to the chosen answer as 1 (never), 2 (rarely), 3  
53 (sometimes), 4 (often). Two positively worded questions, reverse marking from others (“I am  
54 an outgoing person,” and “I can find companionship when I want it”). The total score ranges  
55 from 8-32. A higher score suggested a higher degree of loneliness. We segmented the  
56  
57  
58  
59  
60

1  
2  
3 cumulative score into four groups, no loneliness, 8-16; mild loneliness, 7-21; moderate  
4 loneliness, 22-16; severe loneliness, 27 or above.

### 7 *Patient health questionnaire-9 (PHQ-9)*

9 The PHQ-9 questionnaire has nine different questions for assessing the depressive symptoms  
10 of respondents. The total score ranges from 0 to 27 points where each question has 0-3 points  
11 based on the answers (0, not at all; 1, several days; 2, half of the days; 3, nearly every day).  
12 Below 10 points indicates no depressive symptoms. Ten or higher points indicate the presence  
13 of depressive symptoms.<sup>28</sup> To determine the gravity of depression, we divided the total score  
14 into four levels. The cumulative scores <10, 10-15, 16-21, and 22-27 for no depression, mild  
15 depression, moderate depression, and severe depression, respectively.

### 22 *Generalized anxiety disorder 7-item (GAD-7) scale*

24 The GAD-7 questionnaire consists of seven basic questions for evaluation of anxiety disorder  
25 of respondents. The total score ranges from 0-21. Each question has four different scores  
26 according to responses like 0 (not at all), 1 (several days), 2 (more than half the days), and 3  
27 (nearly every day). A higher score indicates a high degree of anxiety, divided into four different  
28 segments where <5, 5-9, 10-14, and 15-21 indicating no, mild, moderate, and severe anxiety,  
29 respectively.<sup>29</sup>

### 35 *Pittsburgh sleep quality index (PSQI)*

37 We used the PSQI questionnaire to determine the sleep quality of respondents over one month.  
38 This questionnaire consists of nineteen specific questions in seven different domains: (1) sleep  
39 quality (1 question), (2) sleep latency (2 questions), (3) sleep duration (1 question), (4) sleep  
40 efficiency (3 questions), (5) sleep disturbance (9 questions), (6) sleep medication (1 question),  
41 and (7) daily dysfunction (2 questions). Each domain score ranging from 0-3. The cumulative  
42 score of seven domains ranging from 0-21. The higher total scores represent the poor sleep  
43 quality, determining the severity. A total PSQI score below five shows no sleep problem. A  
44 cumulative score of five or more indicates poor sleep. Respondents having a PSQI scores  
45 greater than ten are considered bad sleepers.<sup>30</sup>

### 54 *Statistical analysis*

56 We performed statistical analysis using Microsoft Excel 2016 and statistical packages for social  
57 sciences (IBM SPSS, version 25.0). We used Microsoft Excel for data editing, sorting, coding,  
58 classification, and tabulation. Then we imported the excel file into IBM SPSS software. We  
59



1  
2  
3 used descriptive statistics to analyze the characteristics of the respondents. We applied a chi-  
4 square test to observe the differences in loneliness status (yes or no) with or without depression,  
5 anxiety, or sleep problem among the respondents. We assessed the correlations between risk  
6 factors and psychometric measures (loneliness, depression, anxiety, and sleep quality) using  
7 binary logistic regression analysis with a 95% confidence interval. Statistically significant  
8 results were considered at  $p < 0.05$ .  
9

#### 14 *Patient and public involvement*

16 Patients and the public were not involved in this study.

## 19 **RESULTS**

21 We presented the descriptive statistics for all variables of the respondents in [Table 1](#). Among  
22 the 672 respondents, males and females were 57% and 43%, respectively. Half of the  
23 respondents were above 30 years of age. Among all the respondents, 65% had normal BMI.  
24 About two-thirds of respondents completed their higher secondary education. The married,  
25 non-smoker, medium economic class respondents were 57%, 74%, and 46%, respectively. We  
26 observed two-third of the total respondents are living with family (64%) in a rural area (62%).  
27

28 The estimations of loneliness, depression, anxiety, and sleep problems were 71%, 38%, 64%,  
29 and 73%, respectively ([Figure 1](#)). The proportions of respondents having loneliness were  
30 higher in (i) people living without family members vs. with family members (77% vs. 68%,  
31  $p=0.009$ ), (ii) people with vs. without much depression (89% vs. 60%,  $p < 0.001$ ), anxiety (85%  
32 vs. 46%,  $p < 0.001$ ), and sleep problem (79% vs. 49%,  $p < 0.001$ ), respectively. The sizes of  
33 respondents with depression were higher in (i) females vs. males (48% vs. 31%,  $p < 0.001$ ),  
34 (ii) unemployed vs. service (44% vs. 30%,  $p=0.006$ ), (iii) non-smoker vs. smoker (42% vs.  
35 28%,  $p=0.002$ ), (iv) people with economic status lower vs. medium (43% vs. 33%,  $p=0.047$ ),  
36 (v) respondents living with family members vs. without family members (42% vs. 33%,  
37  $p=0.020$ ), (vi) people with vs. without much loneliness (48% vs. 14%,  $p < 0.001$ ), anxiety (57%  
38 vs. 5%,  $p < 0.001$ ), and sleep problem (45% vs. 19%,  $p < 0.001$ ), respectively. The extents of  
39 respondents with anxiety were higher in (i) females vs. males (71% vs. 69%,  $p=0.001$ ), (ii)  
40 students vs. service (71% vs. 57%,  $p=0.026$ ), (iii) people with BMI below 18.5 vs. above 25  
41 kg/m<sup>2</sup> (88% vs. 63%,  $p=0.017$ ), (iv) people with vs. without much loneliness (77% vs. 32%,  
42  $p < 0.001$ ), depression (95% vs. 44%,  $p < 0.001$ ), and sleep problem (75% vs. 35%,  $p < 0.001$ ),  
43 respectively. And, finally the proportions of respondents with sleep problem were higher in  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 people with vs. without much loneliness (82% vs. 53%,  $p<0.001$ ), depression (87% vs. 65%,  $p$   
4  $< 0.001$ ), and anxiety (85% vs. 53%,  $p<0.001$ ), respectively.  
5  
6

7 We performed the binary logistic regression analysis to measure the correlations between  
8 dependent and independent variables (Table 2). Respondents living with family members were  
9 0.46 times less likely than the respondents living away from family members to have loneliness  
10 (OR=0.46; 95% CI=0.28-0.75,  $p=0.002$ ). Respondents having business occupations were 1.76  
11 times more likely than the unemployed group to have depression (OR=1.76; 95% CI=1.04–  
12 2.99,  $p=0.036$ ). Respondents from the medium economic class were 1.62 times more likely  
13 than those who were from high economic impression to have depression (OR=1.62; 95%  
14 CI=1.04–2.25,  $p=0.033$ ). Urban respondents were 0.66 times less likely to have depression than  
15 rural respondents (OR=0.66; 95% CI=0.43-0.99,  $p=0.044$ ). Respondents living with family  
16 members were 1.80 times more likely to have depression than the respondents living away  
17 from the family members (OR=1.80; 95% CI=1.13-2.87,  $p=0.014$ ). The probabilities of having  
18 loneliness were 2.96 times higher in respondents who already have depression (OR=2.96; 95%  
19 CI=1.73-5.05,  $p<0.001$ ), 3.95 times higher in respondents with anxiety (OR=3.95; 95%  
20 CI=2.51-6.21,  $p<0.001$ ), and 2.64 times higher in respondents suffering from sleep problem  
21 (OR=2.64; 95% CI=1.72-4.05,  $p<0.001$ ), respectively. The chances of having depression were  
22 18.51 times higher in respondents with anxiety (OR=18.51; 95% CI=9.52-35.71,  $p<0.001$ ), and  
23 1.36 times higher in respondents suffering from sleep problem (OR=1.36; 95% CI=0.81-2.35,  
24  $p<0.001$ ), respectively. The likelihoods of having anxiety were 3.60 times higher in  
25 respondents suffering from sleep problems (OR=3.60; 95% CI=2.24-5.65,  $p<0.001$ ).  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

## 40 DISCUSSION

41  
42  
43 This study examines the impact of the COVID-19 pandemic on the mental health of the general  
44 Bangladeshi population. We observed loneliness, depression, anxiety, and sleep problems  
45 among the general population were 71%, 38%, 64%, and 73%, respectively (Figure 1). We  
46 observed the associated factors behind the mental health issues are female gender, low  
47 economic status, students or unemployed, and living without a family. The present finding  
48 reporting a much higher rate of loneliness, depression, anxiety, sleep problems among  
49 Bangladeshi people during the COVID-19 pandemic. Consistent with our study findings, a  
50 cross-sectional epidemiological study reported the higher prevalence of stress (73.4%),  
51 depression (50.7%), anxiety (44.7%), and sleep problems (36.1%) among Chinese people at  
52 the early stage of the outbreak.<sup>31</sup> The higher rates of mental health problems in the present  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 study are consistent with the previous SARS and MERS outbreaks.<sup>8,9</sup> At the early time, people  
4 had little information about the virus, preventive measures and treatment procedures, fatality  
5 rate, etc. The little information and uncertainty about the COVID-19 might contribute to this  
6 higher rate of mental health problems.  
7  
8  
9

10 The present study found that about three in every four people in Bangladesh suffering from  
11 loneliness at any level of gravity during the lockdown period. Among them, 39% were suffering  
12 from moderate to severe loneliness problems. People living without family during lockdown  
13 are more prone to develop loneliness than people living with family. The previous report  
14 suggested that social isolation during the COVID-19 pandemic was a risk factor for  
15 loneliness.<sup>32</sup> People with considerable loneliness are prone to develop other mental health  
16 problems, low well-being, and suicidal behavior.<sup>33,34</sup> Following the current findings, one study  
17 conducted in the United Kingdom and observed 36% of the respondents feel lonely during the  
18 COVID-19 pandemic that is higher than any previous times.<sup>35</sup> Another two studies also  
19 reported higher loneliness scores among general people during COVID-19 than past times.<sup>36,37</sup>  
20 We observed the high prevalence of loneliness among the people living without their family  
21 members. In agreement with the present finding, a recent study reported the increased  
22 loneliness levels among females, young people, singles, unemployed, and those who have other  
23 psychiatric illnesses.<sup>38</sup> Public communication regarding social distancing and mental well-  
24 being involving psychologists, social scientists, and mental health specialists can reduce the  
25 burden of loneliness.<sup>39</sup>  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Among the respondents, 38% experienced depressive symptoms, including mild (24%),  
40 moderate (11%), and severe (3%) levels. We also observed the higher prevalence of depressive  
41 symptoms among females, low economic class people, unemployed people, students, and  
42 people living without a family. Several previous studies also reported a higher rate of  
43 depressive symptoms among females than males during the COVID-19 pandemic.<sup>40-43</sup>  
44 Comparing with the present study results, the increased rate of depression reported in several  
45 recent studies in Spain, China, and Hong Kong due to the COVID-19 pandemic.<sup>44-46</sup> However,  
46 some previous studies reported depression symptoms among the general Chinese population  
47 was 16.5% and 11.4% among Japanese people.<sup>47,48</sup> These inconsistencies might be the result  
48 of developed socioeconomic status and healthcare facilities. Among the study participants,  
49 64% reported anxiety symptoms where mid, moderate, and severe cases were 30%, 17%, and  
50 17%, respectively. Similar to our finding, a study reported the prevalence rate of anxiety  
51 disorder was 28.8% ranging from moderate to severe symptoms in China.<sup>49</sup> Also, a high  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 prevalence of anxiety disorder was observed among respondents from many countries in the  
4 COVID-19 pandemic than previous times.<sup>4</sup> 40 The prevalence rate of anxiety among  
5 Bangladeshi students was 71% in the present study. The final year students of different  
6 education levels might contribute to this high rate due to their uncertain exams and the job  
7 market. We observed, 73% of general people were suffering from sleep problems during the  
8 COVID-19 pandemic. Among them, 50%, 18%, and 5% reported mild, moderate, and severe  
9 sleep problems. The present study also demonstrated that respondents having loneliness,  
10 depression, or anxiety are more likely to have sleep problems than healthy individuals. Many  
11 recent studies reported similar findings. A previous study in China observed 29.2% insomnia,  
12 27.9% depressive symptoms, and 31.6% anxiety disorder among general people during the  
13 COVID-19 pandemic.<sup>50</sup> Similarly, another study observed 42.2% sleep disturbances, 17.4%  
14 moderate or severe insomnia, 24.7% depressive symptoms, and 23.2% anxiety symptoms  
15 among Italian people.<sup>51</sup>

16  
17 To prevent the quick spread of COVID-19 infection, the Bangladesh government postponed  
18 all educational institutions since March 18, 2020. A countrywide lockdown was imposed to  
19 limit public movement since March 26, 2020, and inspired people to stay home.<sup>17</sup> The  
20 countrywide movement restriction and stay home order greatly impacted the economy in the  
21 markets, offices, business organizations, and transport systems.<sup>52</sup> <sup>53</sup> Most of the Bangladeshi  
22 population depends on a regular income. They were uncertain about returning to the workplace  
23 due to indefinite lockdown. This situation created a confounding impact on their mental health  
24 status.<sup>54</sup> In Bangladesh, we observed several reported suicide cases during the COVID-19  
25 pandemic for getting infected, economic loss, social security, job security, and the emotional  
26 breakdown of marginalized wage-earners.<sup>55</sup> <sup>56</sup> An Indian case study suggested that COVID-19  
27 may significantly impact the mental health status and influence suicidal ideation, suicide  
28 attempts among the affected people; other comorbid diseases may aggravate the situation.<sup>57</sup>  
29 We found that 44% of unemployed and 30% of service-holders were suffering from depression.  
30 About 43% of people belonging to the lower economic class are facing depression during the  
31 lockdown period.

32  
33 The prevalence rates of mental health problems before the COVID-19 period varied from 6.5%  
34 to 31.0% among adults in Bangladesh.<sup>58</sup> The significant increase of prevalence rates of mental  
35 health issues during the COVID-19 pandemic (ranging from 38% to 73%) in the present study  
36 is a concern. Therefore, the National Institute of Mental Health of Bangladesh has announced  
37 some recommendations. These are psychotropics, avoid COVID-19 news multiple times in a

1  
2  
3 day or news scrolling, one or two authentic sources of information, less social media, simple  
4 relaxation technique like breathing exercise, etc., regarding the management of mental health  
5 during the COVID-19 pandemic.<sup>59</sup> In agreement with the above suggestions, several studies  
6 also recommended online response, counseling, social support, training on mental health for  
7 the patients, health care professionals, public service holders, youth, students, elderly  
8 populations to manage mental health problems.<sup>60-62</sup> Worldwide, the COVID-19 pandemic  
9 unveiled the actual setup of unpreparedness of the healthcare systems and the scarcity of  
10 resources (PPE, testing kits, etc.) to combat the situation.<sup>63</sup>  
11  
12  
13  
14  
15  
16  
17

### 18 **Strengths and limitations of this study**

19  
20 The present study has some limitations. First, online self-reporting surveys might have multiple  
21 biases and are not representative of those without internet facilities. Second, this cross-sectional  
22 study is not able to measure the impact of these mental illnesses over time. Third, we didn't  
23 assess the altered life-style of the respondents during the COVID-19 pandemic. Besides the  
24 limitations, the present study has some advantages. Firstly, this is the first-ever study in  
25 Bangladesh assessing the four major mental health issues during the COVID-19 pandemic.  
26 This study provides an idea about the mental health of the Bangladeshi population during the  
27 COVID-19 pandemic.  
28  
29  
30  
31  
32  
33

### 34 **Future research**

35  
36 This study highlights the need for mental health assessment and proper management of these  
37 issues during the COVID-19 pandemic and future research among healthcare professionals to  
38 explore their actual mental health status in this pandemic situation.  
39  
40  
41

### 42 **CONCLUSION**

43  
44 In summary, our findings reflect the gravity of mental health problems during the COVID-19  
45 pandemic. A large portion of general people in Bangladesh affected mentally with different  
46 levels of severity. We suggest intensive mental healthcare services for the Bangladeshi people.  
47 Therefore, integrated government, non-government, and community activities can ensure  
48 individual and collective mental health. Besides mental health support, social security and  
49 economic stability should be the top priority to grow the confidence among general people.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Declarations

## Acknowledgments

All the authors are thankful to the participants for their cooperation to conduct this study.

## Author contributions

RD conceived the idea, designed the study, collected the data, and prepared the initial draft of the manuscript. MRH and SD contributed to the development of the study design, collected the data, and edited the manuscript. MRI conceived and designed the study, analyzed and interpreted the data, reviewed the manuscript for important intellectual content, and supervised the whole work. All the authors approved the final version of the manuscript for this submission.

## Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

## Ethical statement

Our study did not require an ethical board approval because it did not contain human or animal trials.

## Competing interests

The authors do not have any competing interests to declare.

## Patient consent for publication

Not applicable.

## Data availability statement

Data supporting the present study findings are within this article. All the relevant data and information can be obtained from the corresponding author upon reasonable request.

## Table and Figure Legends

**Table1** Distribution of variables and their association with different mental health problems among the respondents.

p-values are significant at 95% confidence interval ( $p < 0.05$ ). Significant p-values are shown in bold.

BMI, Body Mass Index; df, Degrees of Freedom; N, Number.

**Table 2** Regression analysis of variables by mental health problems among respondents.

p-values are significant at 95% confidence interval ( $p < 0.05$ ). Significant p-values are shown in bold.

BMI, Body Mass Index; CI, Confidence Interval; N, Number; OR, Odds Ratio.

**Figure 1** Mental health problems among the respondents based on their distribution and gravity.

## Reference

1. Talevi D, Socci V, Carai M, et al. Mental health outcomes of the COVID-19 pandemic. *Rivista di Psichiatria*. 2020;55(3):137-144.
2. WHO COVID-19 Weekly Epidemiological Update - 29 December 2020. Available: <https://www.who.int/publications/m/item/weekly-epidemiological-update---29-december-2020> [accessed February 6, 2021].
3. WHO Coronavirus Disease (COVID-19) Dashboard. Available: [https://covid19.who.int/?gclid=Cj0KCQiAmfmABhCHARIsACwPRADLrk2VaTyZeSFy6kf8pjyrLYNDLGHR0c8z4nYnqSyLggvDD63O9eAaAiK0EALw\\_wcB](https://covid19.who.int/?gclid=Cj0KCQiAmfmABhCHARIsACwPRADLrk2VaTyZeSFy6kf8pjyrLYNDLGHR0c8z4nYnqSyLggvDD63O9eAaAiK0EALw_wcB) [accessed February 6, 2021].
4. Chen Y, Zhou H, Zhou Y, Zhou F. Prevalence of self-reported depression and anxiety among pediatric medical staff members during the COVID-19 outbreak in Guiyang, China. *Psychiatry Res*. 2020;288:113005.
5. Duan L, Zhu G. Psychological interventions for people affected by the COVID-19 epidemic. *Lancet Psychiatry*. 2020;7(4):300-302.
6. Madabhavi I, Sarkar M, Kadakol N. COVID-19: a review. *Monaldi Arch Chest Dis* 2020;90(2).
7. Zhang Y, Ma ZF. Impact of the COVID-19 Pandemic on Mental Health and Quality of Life among Local Residents in Liaoning Province, China: A Cross-Sectional Study. *Int J Environ Res Public Health*. 2020;17(7):2381.
8. Lau JT, Griffiths S, Choi KC, Tsui HY. Avoidance behaviors and negative psychological responses in the general population in the initial stage of the H1N1 pandemic in Hong Kong. *BMC Infect Dis*. 2010;10:139.
9. Jeong H, Yim HW, Song YJ, Ki M, Min JA, Cho J, Chae JH. Mental health status of people isolated due to Middle East Respiratory Syndrome. *Epidemiol Health*. 2016;38:e2016048.
10. Maunder RG, Lancee WJ, Balderson KE, et al. Long-term psychological and occupational effects of providing hospital healthcare during SARS outbreak. *Emerg Infect Dis*. 2006;12(12):1924-1932.
11. North CS, Pfefferbaum B, Vythilingam M, et al. Exposure to bioterrorism and mental health response among staff on Capitol Hill. *Biosecur Bioterror*. 2009;7(4):379-388.

12. Xiang YT, Yang Y, Li W, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry*. 2020;7(3):228-229.
13. Central intelligence agency. The world factbook, 2020. Available: [https://www.cia.gov/library/publications/resources/the-world-factbook/geos/print\\_bg.html](https://www.cia.gov/library/publications/resources/the-world-factbook/geos/print_bg.html) [Accessed 01 September 2020].
14. Streatfield PK, Karar ZA. Population challenges for Bangladesh in the coming decades. *J Health Popul Nutr*. 2008;26(3):261-272.
15. Paul R. Bangladesh Confirms its First Three Cases of Coronavirus, 2020. Available: <https://www.reuters.com/article/us-healthcoronavirus-bangladesh-idUSKBN20V0FS> [Accessed 14 April 2020].
16. WHO Bangladesh COVID-19 Morbidity and Mortality Weekly Update - 27 - December 2020 (MMWU). Available: [https://cdn.who.int/media/docs/default-source/searo/bangladesh/covid-19-who-bangladesh-situation-reports/who\\_covid-19-update\\_44\\_20201227.pdf?sfvrsn=bffee8db\\_7](https://cdn.who.int/media/docs/default-source/searo/bangladesh/covid-19-who-bangladesh-situation-reports/who_covid-19-update_44_20201227.pdf?sfvrsn=bffee8db_7) [accessed February 6, 2021].
17. The Daily Star. Man left out of relief net dies by suicide in Chattogram, 2020. Available: <https://www.thedailystar.net/coronavirus-deadly-new-threat/news/man-left-out-relief-net-dies-suicide-chattogram-1893409> [Accessed 16 April 2020].
18. The Kaler Kantho. Day labourer commits suicide in Naldanga (In Bangla), 2020. Available: <https://www.kalerkantho.com/home/printnews/898402/2020-04-13> [Accessed 13 April 2020].
19. The Jugantor. The young woman committed suicide after losing her job in Corona left posting on Facebook (In Bangla), 2020. Retrieved July 11, 2020, Available: <https://rb.gy/pmssav> [Accessed 11 July 2020].
20. Silva Junior FJGD, Sales JCES, Monteiro CFS, Costa APC, Campos LRB, Miranda PIG, Monteiro TAS, Lima RAG, Lopes-Junior LC. Impact of COVID-19 pandemic on mental health of young people and adults: a systematic review protocol of observational studies. *BMJ Open*. 2020;10(7):e039426.
21. Lopes-Júnior LC, Bomfim E, Silveira DSCD, Pessanha RM, Schuab SIPC, Lima RAG. Effectiveness of mass testing for control of COVID-19: a systematic review protocol. *BMJ Open*. 2020;10(8):e040413.
22. Every-Palmer S, Jenkins M, Gendall P, Hoek J, Beaglehole B, Bell C, Williman J, Rapsey C, Stanley J. Psychological distress, anxiety, family violence, suicidality, and wellbeing in New Zealand during the COVID-19 lockdown: A cross-sectional study. *PLoS One*. 2020;15(11):e0241658.
23. Chen Q, Liang M, Li Y, Guo J, Fei D, Wang L, He L, Sheng C, Cai Y, Li X, Wang J, Zhang Z. Mental health care for medical staff in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020 Apr;7(4):e15-e16. doi: 10.1016/S2215-0366(20)30078-X. Epub 2020 Feb 19. Erratum in: *Lancet Psychiatry*. 2020;7(5):e27.
24. Yang Y, Li W, Zhang Q, Zhang L, Cheung T, Xiang YT. Mental health services for older adults in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020;7(4):e19.



- 1
- 2
- 3
- 4 25. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, Rubin GJ. The psychological impact of quarantine and how to reduce it: rapid review of the
- 5 evidence. *Lancet*. 2020;395(10227):912-920.
- 6
- 7
- 8 26. Beaton DE, Bombardier C, Guillemin F, Ferraz MB. Guidelines for the process of cross-
- 9 cultural adaptation of self-report measures. *Spine*. 2000; 25:3186–91.
- 10
- 11 27. Hays RD, DiMatteo MR. A short-form measure of loneliness. *J Pers Assess*.
- 12 1987;51(1):69-81.
- 13
- 14 28. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity
- 15 measure. *J Gen Intern Med*. 2001;16(9):606-613.
- 16
- 17 29. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized
- 18 anxiety disorder: the GAD-7. *Arch Intern Med*. 2006;166(10):1092-1097.
- 19
- 20 30. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta psychiatrica*
- 21 *scandinavica*. 1983;67(6):361-370.
- 22
- 23 31. Liu S, Yang L, Zhang C, Xiang YT, Liu Z, Hu S, Zhang B. Online mental health
- 24 services in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020;7(4):e17-e18.
- 25
- 26 32. Dahlberg L. Loneliness during the COVID-19 pandemic. *Aging Ment Health*.
- 27 2021;25:1-4.
- 28
- 29 33. Leigh-Hunt N, Bagguley D, Bash K, Turner V, Turnbull S, Valtorta N, Caan W. An
- 30 overview of systematic reviews on the public health consequences of social isolation and
- 31 loneliness. *Public Health*. 2017;152:157-171.
- 32
- 33 34. Rico-Uribe LA, Caballero FF, Martín-María N, Cabello M, Ayuso-Mateos JL, Miret M.
- 34 Association of loneliness with all-cause mortality: A meta-analysis. *PLoS One*.
- 35 2018;13(1):e0190033.
- 36
- 37 35. Li LZ, Wang S. Prevalence and predictors of general psychiatric disorders and loneliness
- 38 during COVID-19 in the United Kingdom [published online ahead of print, 2020 Jun
- 39 30]. *Psychiatry Res*. 2020;291:113267.
- 40
- 41 35. Luchetti M, Lee JH, Aschwanden D, et al. The trajectory of loneliness in response to
- 42 COVID-19 [published online ahead of print, 2020 Jun 22]. *Am Psychol*.
- 43 2020;10.1037/amp0000690.
- 44
- 45 37. Killgore WDS, Cloonan SA, Taylor EC, Dailey NS. Loneliness: A signature mental
- 46 health concern in the era of COVID-19. *Psychiatry Res*. 2020;290:113117.
- 47
- 48 38. Hoffart A, Johnson SU, Ebrahimi OV. Loneliness and Social Distancing During the
- 49 COVID-19 Pandemic: Risk Factors and Associations With Psychopathology. *Front*
- 50 *Psychiatry*. 2020;11:589127.
- 51
- 52 39. Popa E. Loneliness and negative effects on mental health as trade-offs of the policy
- 53 response to COVID-19. *Hist Philos Life Sci*. 2021;43(1):15.
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3
- 4 40. Zhou SJ, Zhang LG, Wang LL, et al. Prevalence and socio-demographic correlates of
- 5 psychological health problems in Chinese adolescents during the outbreak of COVID-19.
- 6 *Eur Child Adolesc Psychiatry*. 2020;29(6):749-758.
- 7
- 8 41. Islam MR, Ali S, Karmoker JR, et al. Evaluation of serum amino acids and non-
- 9 enzymatic antioxidants in drug-naïve first-episode major depressive disorder. *BMC*
- 10 *Psychiatry*. 2020;20(1):333.
- 11
- 12 42. Li G, Miao J, Wang H, et al. Psychological impact on women health workers involved in
- 13 COVID-19 outbreak in Wuhan: a cross-sectional study. *J Neurol Neurosurg Psychiatry*.
- 14 2020;91(8):895-897.
- 15
- 16 43. Kang L, Ma S, Chen M, et al. Impact on mental health and perceptions of psychological
- 17 care among medical and nursing staff in Wuhan during the 2019 novel coronavirus
- 18 disease outbreak: A cross-sectional study. *Brain Behav Immun*. 2020;87:11-17.
- 19
- 20 44. González-Sanguino C, Ausín B, Castellanos MÁ, et al. Mental health consequences
- 21 during the initial stage of the 2020 Coronavirus pandemic (COVID-19) in Spain. *Brain*
- 22 *Behav Immun*. 2020;87:172-176.
- 23
- 24 45. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality
- 25 during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry*
- 26 *Res*. 2020;288:112954.
- 27
- 28 46. Choi EPH, Hui BPH, Wan EYF. Depression and Anxiety in Hong Kong during COVID-
- 29 19. *Int J Environ Res Public Health*. 2020;17(10):3740.
- 30
- 31 47. Wang C, Pan R, Wan X, Tan Y, Xu L, Mc Intyre RS, Choo FN, Tran B, Ho R, Sharma
- 32 VK. A longitudinal study on the mental health of general population during the COVID-
- 33 19 epidemic in China. *Brain Behav Immun*. 2020;87:40–48.
- 34
- 35 48. Ueda M, Stickley A, Sueki H, Matsubayashi T 2020. Mental health status of the general
- 36 population during the COVID-19 pandemic: a cross-sectional national survey in Japan.
- 37 *medRxiv*. [accessed Feb 7, 2021].
- 38 <https://www.medrxiv.org/content/10.1101/2020.04.28.20082453v3>
- 39
- 40 49. Liu X, Luo WT, Li Y, Li CN, Hong ZS, Chen HL, Xiao F, Xia JY. Psychological status
- 41 and behavior changes of the public during the COVID-19 epidemic in China. *Infect Dis*
- 42 *Poverty*. 2020;9(1):58.
- 43
- 44 50. Shi L, Lu ZA, Que JY, et al. Prevalence of and Risk Factors Associated With Mental
- 45 Health Symptoms Among the General Population in China During the Coronavirus
- 46 Disease 2019 Pandemic. *JAMA Netw Open*. 2020;3(7):e2014053.
- 47
- 48 51. Gualano MR, Lo Moro G, Voglino G, Bert F, Siliquini R. Effects of Covid-19
- 49 Lockdown on Mental Health and Sleep Disturbances in Italy. *Int J Environ Res Public*
- 50 *Health*. 2020;17(13):4779.
- 51
- 52 52. The Bangla Tribune. The patient committed suicide by jumping from the roof as he was
- 53 infected with corona (In Bangla), 2020. Available: <https://rb.gy/jfjogjx> [Accessed 11 June
- 54 2020].
- 55
- 56
- 57
- 58
- 59
- 60

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
53. The Somoy News. Patient commits suicide without getting treatment in hospital (In Bangla), 2020. Available: <https://www.somoynews.tv/pages/details/214750> [Accessed 20 May 2020].
54. Hossain KA, Roy S, Ullah MM, Kabir R, Arafat SY. COVID-19 and Mental Health Challenges in Bangladesh. *Advances in Medical, Dental and Health Sciences*. 2020;2(3)31-3.
55. The BBC News. Corona virus: COVID-19 patient committed suicide after escaping from the hospital (In Bangla), 2020. Available: <https://www.bbc.com/bengali/news-53125017> [Accessed 21 June 2020].
56. The Desh Rupantor. Corona-infected businessman commits suicide (In Bangla), 2020. Available: <https://www.deshrupantor.com/mofossol/2020/07/07/230368> [Accessed 07 July 2020].
57. Hossain MM, Tasnim S, Sultana A, McKyer EL, Ma P. COVID-19 and suicide of an army soldier in India: perspectives on psychosocial epidemiology of suicidal behavior. *SocArXiv*. 2020.
58. Hossain MD, Ahmed HU, Chowdhury WA, Niessen LW, Alam DS. Mental disorders in Bangladesh: a systematic review. *BMC Psychiatry*. 2014;14:216.
59. National Institute of Mental Health (MIMH). Recommendations from Sub Committee for Mental Health, 2020. Available: [https://mofa.portal.gov.bd/sites/default/files/files/mofa.portal.gov.bd/page/ad1f289c\\_47cf\\_4f6c\\_8dee\\_887957be3176/Guideline%20for%20Mental%20Health%20Issue-COVID-19-%20DGHS.pdf](https://mofa.portal.gov.bd/sites/default/files/files/mofa.portal.gov.bd/page/ad1f289c_47cf_4f6c_8dee_887957be3176/Guideline%20for%20Mental%20Health%20Issue-COVID-19-%20DGHS.pdf) [Accessed 01 September, 2020]
60. Wang C, Pan R, Wan X, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health*. 2020;17(5):1729.
61. Yang Y, Li W, Zhang Q, Zhang L, Cheung T, Xiang YT. Mental health services for older adults in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020;7(4):e19.
62. Armitage R, Nellums LB. The COVID-19 response must be disability inclusive. *Lancet Public Health*. 2020;5(5):e257.
63. Pfefferbaum B, North CS. Mental Health and the Covid-19 Pandemic. *N Engl J Med*. 2020;383(6):510-512.

**Table 1** Distribution of variables and their association with different mental health problems among the respondents.

	Total N=672		Loneliness N=478				Depression N=256				Generalized anxiety N=429				Sleep problem N=494								
	n	%	Yes n	% %	$\chi^2$	df	p-value	Yes n	% %	$\chi^2$	df	p-value	Yes n	% %	$\chi^2$	df	p-value	Yes n	% %	$\chi^2$	df	p-value	
<b>Age</b>																							
15 to 30 years	344	51	236	69	2.19	1	0.139	119	35	3.67	1	0.056	213	62	1.12	1	0.289	242	70	3.62	1	0.051	
Above 30 years	328	49	242	74				137	42				216	66				252	77				
<b>Gender</b>																							
Male	381	57	278	73	1.44	1	0.230	117	31	20.35	1	<0.001	223	59	10.74	1	0.001	271	71	2.57	1	0.109	
Female	291	43	200	69				139	48				206	71				233	80				
<b>BMI (kg/m<sup>2</sup>)</b>																							
Below 18.5	32	5	24	75	0.25	2	0.880	18	56	4.74	2	0.093	28	88	8.16	2	0.017	26	81	1.08	2	0.584	
18.5–25	435	65	308	71				163	37				272	63				317	73				
Above 25	205	31	146	71				75	37				129	63				151	74				
<b>Marital status</b>																							
Unmarried	289	43	210	73	0.58	1	0.446	107	37	0.25	1	0.619	187	65	0.16	1	0.685	207	72	0.93	1	0.336	
Married	383	57	268	70				149	39				242	63				287	75				
<b>Education</b>																							
Illiterate	26	4	21	81	7.53	4	0.110	15	58	6.00	4	0.199	18	69	7.39	4	0.117	19	73	0.71	4	0.849	
Primary	85	13	59	69				35	41				61	72				65	76				
Secondary	61	9	48	79				23	38				44	72				44	72				
Higher Secondary	410	61	279	68				146	36				246	60				302	74				
Graduate/above	90	13	71	79				37	41				60	67				64	71				
<b>Occupation</b>																							
Service	219	33	153	70	3.75	3	0.290	66	30	12.45	3	0.006	125	57	9.28	3	0.026	158	72	0.56	3	0.906	
Business	28	4	20	71				7	25				17	61				20	71				
Student	208	31	158	76				88	42				148	71				153	74				
Unemployed	217	32	147	68				95	44				139	64				163	75				

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

Table 1 (Continued)

<b>Economic status</b>																							
Low	245	36	179	73	0.75	2	0.688	106	43	6.11	2	<b>0.047</b>	163	67	1.39	2	0.499	177	72	2.60	2	0.273	
Medium	310	46	218	70				103	33				195	63				224	72				
High	117	17	81	69				47	40				71	61				93	79				
<b>Residence</b>																							
Urban	255	38	184	72	0.21	1	0.646	109	43	3.79	1	0.052	164	64	0.04	1	0.841	192	75	0.67	1	0.413	
Rural	417	62	194	47				147	35				265	64				302	72				
<b>Living status</b>																							
With family	430	64	291	68	11.62	3	<b>0.009</b>	179	42	9.86	3	<b>0.020</b>	274	64	1.56	3	0.665	323	75	4.82	3	0.185	
Without family	242	36	187	77				77	32				155	64				171	71				
<b>Smoking habit</b>																							
Smoker	177	26	135	76	3.09	1	0.079	50	28	9.88	1	<b>0.002</b>	105	59	2.12	1	0.145	129	73	0.05	1	0.825	
Non-smoker	495	74	343	69				206	42				324	65				365	74				
<b>Loneliness</b>																							
Yes	478	71	478	100	-	-	-	228	48	64.75	1	<b>&lt;0.001</b>	366	77	116.23	1	<b>&lt;0.001</b>	391	82	58.40	1	<b>&lt;0.001</b>	
No	194	29	0	0				28	14				63	32				103	53				
<b>Depression</b>																							
Yes	256	38	228	89	64.75	1	<b>&lt;0.001</b>	256	100	-	-	-	244	95	177.45	1	<b>&lt;0.001</b>	222	87	37.04	1	<b>&lt;0.001</b>	
No	416	62	250	60				0	0				185	44				272	65				
<b>Generalized anxiety</b>																							
Yes	429	64	366	85	116.23	1	<b>&lt;0.001</b>	244	57	177.45	1	<b>&lt;0.001</b>	429	100	-	-	-	366	85	84.86	1	<b>&lt;0.001</b>	
No	243	36	112	46				12	5				0	0				128	53				
<b>Sleep problem</b>																							
Yes	494	75	391	79	58.40	1	<b>&lt;0.001</b>	222	45	37.04	1	<b>&lt;0.001</b>	366	75	84.88	1	<b>&lt;0.001</b>	494	100	-	-	-	
No	178	25	87	49				34	19				63	35				0	0				

p-values are significant at 95% confidence interval (p<0.05). Significant p-values are shown in bold.

BMI, Body Mass Index; df, Degrees of Freedom; N, Number.

**Table 2** Regression analysis of variables by mental health problems among respondents.

	Loneliness N=478			Depression N=256			Generalized anxiety N=429			Sleep problem N=494		
	OR	95% CI	p-value	OR	95% CI	p-value	OR	95% CI	p-value	OR	95% CI	p-value
<b>Age</b>												
15 to 30 years	0.606	0.359-1.023	0.061	0.653	0.386-1.105	0.112	0.857	0.490-1.501	0.590	0.834	0.499-1.394	0.489
Above 30 years	1			1			1			1		
<b>Gender</b>												
Male	0.666	0.411-1.080	0.099	1.624	1.000-2.637	<b>0.050</b>	1.453	0.869-2.431	0.154	1.036	0.648-1.658	0.882
Female	1			1			1			1		
<b>BMI (kg/m<sup>2</sup>)</b>												
Below 18.5	1.574	0.557-4.443	0.392	0.889	0.346-2.282	0.806	0.270	0.072-1.011	0.052	0.818	0.289-2.319	0.706
18.5-25	1.244	0.399-3.877	0.706	0.832	0.287-2.410	0.735	0.347	0.085-1.411	0.139	0.877	0.281-2.737	0.821
Above 25	1			1			1			1		
<b>Marital status</b>												
Unmarried	0.902	0.508-1.600	0.723	0.845	0.463-1.542	0.584	1.097	0.599-2.011	0.763	1.009	0.578-1.764	0.974
Married	1			1			1			1		
<b>Education</b>												
Illiterate	0.498	0.211-1.172	0.111	1.076	0.476-2.430	0.861	0.511	0.215-1.211	0.127	1.346	0.617-2.935	0.455
Primary	1.365	0.491-3.794	0.551	1.545	0.620-3.851	0.351	0.566	0.210-1.525	0.260	0.855	0.350-2.090	0.732
Secondary	1.123	0.293-4.309	0.866	1.813	0.507-6.485	0.360	0.244	0.059-1.006	0.051	0.829	0.249-2.763	0.761
Higher Secondary	0.518	0.204-1.314	0.166	1.006	0.430-2.351	0.989	0.788	0.311-2.000	0.616	1.396	0.589-3.310	0.449
Graduate/above	1			1			1			1		
<b>Occupation</b>												
Service	0.889	0.292-2.711	0.836	0.315	0.099-1.009	0.052	0.855	0.296-2.481	0.774	0.894	0.318-2.515	0.831
Business	1.476	0.895-2.434	0.127	1.760	1.037-2.994	0.036	1.300	0.754-2.237	0.344	1.161	0.702-1.923	0.559
Student	0.628	0.264-1.962	0.253	0.752	0.363-1.052	0.093	0.548	0.362-0.952	0.452	0.524	0.241-1.183	0.263
Unemployed	1			1			1			1		
<b>Economic status</b>												
Low	1.111	0.636-1.941	0.712	1.478	0.832-2.624	0.183	0.562	0.312-1.011	0.054	1.588	0.893-2.824	0.115
Medium	1.133	0.725-1.771	0.583	1.618	1.038-2.252	<b>0.033</b>	0.981	0.613-1.572	0.938	0.852	0.558-1.300	0.458
High	1			1			1			1		

Table 2 (Continued)

<b>Residence</b>												
Urban	1.06	0.697-1.611	0.785	0.655	0.434-0.988	<b>0.044</b>	1.146	0.737-1.781	0.545	0.929	0.618-1.397	0.724
Rural	1			1			1			1		
<b>Living status</b>												
With family	0.45	0.278-0.751	<b>0.002</b>	1.797	1.126-2.867	<b>0.014</b>	0.842	0.511-1.388	0.501	1.344	0.844-2.140	0.213
Without family	1			1			1			1		
<b>Smoking habit</b>												
Smoker	0.63	0.376-1.078	0.093	1.441	0.849-2.447	0.176	1.032	0.611-1.743	0.906	0.925	0.567-1.509	0.755
Non-smoker	1			1			1			1		
<b>Loneliness</b>												
Yes	-	-	-	2.890	1.669-5.000	<b>&lt;0.001</b>	3.952	2.500-6.211	<b>&lt;0.001</b>	2.525	1.647-3.875	<b>&lt;0.001</b>
No	-			1			1			1		
<b>Depression</b>												
Yes	2.95	1.733-5.050	<b>&lt;0.00</b>	-	-	-	18.518	9.615-	<b>&lt;0.001</b>	1.344	0.800-2.257	0.263
No	1			-			1			1		
<b>Generalized anxiety</b>												
Yes	3.95	2.506-6.211	<b>&lt;0.00</b>	18.518	9.523-35.714	<b>&lt;0.001</b>	-	-	-	3.367	2.127-3.519	<b>&lt;0.001</b>
No	1			1			-			1		
<b>Sleep problem</b>												
Yes	2.63	1.718-4.048	<b>&lt;0.00</b>	1.385	0.814-2.352	0.229	3.558	2.237-5.649	<b>&lt;0.001</b>	-	-	-
No	1			1			1			-		

p-values are significant at 95% confidence interval (p<0.05). Significant p-values are shown in bold.

BMI, Body Mass Index; CI, Confidence Interval; N, Number; OR, Odds Ratio.




Figure 1: Mental health problems among the respondents based on their distribution and gravity



**Title: Impact of COVID-19 pandemic on mental health among general Bangladeshi population: A cross-sectional study**

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Section/Paragraph
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	Title (page 1)
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Abstract (page 2)
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Introduction Paragraphs 1-3 (Page 4-5)
Objectives	3	State specific objectives, including any prespecified hypotheses	Introduction Paragraphs 4 (Page 5)
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	Methods Paragraphs 1 (Page 5)
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Methods Paragraphs 1 (Page 5)
Participants	6	(a) Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls	Methods Paragraphs 1 (Page 5)
		(b) For matched studies, give matching criteria and the number of controls per case	N/A
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Methods Paragraphs 1 (Page 5)
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Methods Paragraphs 2-3 (Page 6)
Bias	9	Describe any efforts to address potential sources of bias	Methods Paragraphs 1 (Page 5)
Study size	10	Explain how the study size was arrived at	Methods Paragraphs 1 (Page 5)
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Methods Paragraphs 1 (Page 5)
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Methods Paragraphs 8 (Page 7-8)
		(b) Describe any methods used to examine subgroups and interactions	Methods Paragraphs 8 (Page 7-8)
		(c) Explain how missing data were addressed	Methods Paragraphs 1 (Page 5)
		(d) If applicable, explain how matching of cases and controls was addressed	N/A
		(e) Describe any sensitivity analyses	N/A
<b>Results</b>			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Results Paragraphs 1 (Page 8)
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Results Paragraphs 2 (Page 8)
		(b) Indicate number of participants with missing data for each variable of interest	N/A
Outcome data	15*	Report numbers in each exposure category, or summary measures of exposure	Results Paragraphs 2 (Page 9)
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Results Paragraphs 2 (Page 9) Table 1 and 2
		(b) Report category boundaries when continuous variables were categorized	N/A
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	N/A
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	Discussion Paragraph 1 (page 9-10)
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Strengths and limitations section (page 12)
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Discussion Paragraph 2-4 (page 10-12)
Generalisability	21	Discuss the generalisability (external validity) of the study results	Discussion Paragraph 3-4 (page 11)
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Page 13

\*Give information separately for cases and controls.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at <http://www.strobe-statement.org>.